

Conditional Analysis on \mathbb{R}^d

Patrick Cheridito*

Michael Kupper[†]

Nicolas Vogelpoth[‡]

October 2014[§]

Abstract. This paper provides versions of classical results from linear algebra, real analysis and convex analysis in a free module of finite rank over the ring L^0 of measurable functions on a σ -finite measure space. We study the question whether a submodule is finitely generated and introduce the more general concepts of L^0 -affine sets, L^0 -convex sets, L^0 -convex cones, L^0 -hyperplanes and L^0 -halfspaces. We investigate orthogonal complements, orthogonal decompositions and the existence of orthonormal bases. We also study L^0 -linear, L^0 -affine, L^0 -convex and L^0 -sublinear functions and introduce notions of continuity, differentiability, directional derivatives and subgradients. We use a conditional version of the Bolzano–Weierstrass theorem to show that conditional Cauchy sequences converge and give conditions under which conditional optimization problems have optimal solutions. We prove results on the separation of L^0 -convex sets by L^0 -hyperplanes and study L^0 -convex conjugate functions. We provide a result on the existence of L^0 -subgradients of L^0 -convex functions, prove a conditional version of the Fenchel–Moreau theorem and study conditional inf-convolutions.

Keywords: L^0 -modules, random sets, conditional optimization, L^0 -differentiability, L^0 -convexity, separating L^0 -hyperplanes, L^0 -convex conjugation, L^0 -subgradients.

2010 Mathematics Subject Classification: 13C13, 46A19, 46A22, 60H25

1 Introduction

Let L^0 be the set of all real-valued measurable functions on a σ -finite measure space $(\Omega, \mathcal{F}, \mu)$, where two of them are identified if they agree μ -almost everywhere. The purpose of this paper is to study the set $(L^0)^d$ of all d -dimensional vectors with components in L^0 and functions $f : (L^0)^d \rightarrow L^0$. Its main motivation are applications in the following two special cases:

- If μ is a probability measure, the elements of L^0 are random variables, and subsets $C \subseteq (L^0)^d$ can be understood as random sets in \mathbb{R}^d . A typical function $f : (L^0)^d \rightarrow L^0$ would, for example, be a mapping that conditionally on \mathcal{F} , assigns to every random point $X \in (L^0)^d$ its Euclidean distance to C .
- Let $(\Omega, \mathcal{G}, \mu)$ be the product of a σ -finite measure space $(\mathbb{T}, \mathcal{H}, \nu)$ and a probability space (E, \mathcal{E}, P) . If \mathcal{F} is a sub- σ -algebra of \mathcal{G} , the elements of L^0 are stochastic processes $(X_t)_{t \in \mathbb{T}}$ on (E, \mathcal{E}, P) . A subset $C \subseteq (L^0)^d$ could, for instance, describe the set of admissible strategies in a stochastic control problem, and an optimal strategy could be characterized as the conditional optimizer of an appropriate function $f : (L^0)^d \rightarrow L^0$ over C .

Unless Ω is the union of finitely many atoms, $(L^0)^d$ is an infinite-dimensional vector space over \mathbb{R} . But conditioned on \mathcal{F} , it is only d -dimensional. Or put differently, it is a free module of rank d over the ring L^0 . This allows us to derive conditional analogs of classical results from linear algebra, real analysis and

*Princeton University, Princeton, NJ 08544, USA.

[†]University of Konstanz, Universitätsstraße 10, 78464 Konstanz Germany.

[‡]20 Main Mill, Mumfords Mill, 19-23 Greenwich High Road, SE10 8ND London, UK.

[§]We thank Ramon van Handel, Ying Hu, Asgar Jamneshan, Mitja Stadje and Martin Streckfuß for fruitful discussions and helpful comments.

convex analysis that depend on the fact that \mathbb{R}^d is a finite-dimensional vector space. L^0 -modules have been studied before; see, for instance, Filipović et al. (2009), Kupper and Vogelpoth (2009), Guo (2010), Guo (2011) and the references in these papers. But since we consider free modules of finite rank, we are able to provide stronger results under weaker assumptions, and moreover, do not need Zorn's lemma or the axiom of choice. Our approach differs from standard measurable selection arguments in that we work modulo null-sets with respect to the measure μ and do not use ω -wise arguments. This has the advantage that one never leaves the world of measurable functions. But it only works in situations where a measure μ is given, and the quantities of interest do not depend on μ -null sets.

The results in this paper are theoretical. But they have already been applied several times: in Cheridito and Hu (2011), they were used to describe stochastic constraints and characterize optimal strategies in a dynamic consumption and investment problem. In Cheridito and Stadje (2012) they guaranteed the existence of a conditional subgradient. In Cheridito et al. (2012) they were applied to show existence and uniqueness of economic equilibria in incomplete market models.

The structure of the paper is as follows: In Section 2 we investigate when an L^0 -submodule of $(L^0)^d$ is finitely generated. Then we study conditional orthogonality and introduce L^0 -affine sets, L^0 -convex sets and L^0 -convex cones. It turns out that the notion of σ -stability plays a crucial role. In Section 3 we investigate almost everywhere converging sequences in $(L^0)^d$ and the corresponding notion of closure. We define L^0 -linear and L^0 -affine functions $f : (L^0)^d \rightarrow (L^0)^k$ and show that they are continuous with respect to almost everywhere converging sequences. We also give a conditional version of the Bolzano–Weierstrass theorem and show that conditional Cauchy sequences converge. Moreover, we define L^0 -bounded sets and give a condition for L^0 -convex sets to be L^0 -bounded. In Section 4 we study sequentially semicontinuous and L^0 -convex functions $f : (L^0)^d \rightarrow L^0$ and prove a result which guarantees that a conditional optimization problem has an optimal solution. Section 5 is devoted to L^0 -open sets, interiors and relative interiors. L^0 -open sets form a topology, but they are not complements of sequentially closed sets. In Section 6 we give strong, weak and proper separation results of L^0 -convex sets by L^0 -hyperplanes. Section 7 studies L^0 -convex functions and introduces conditional notions of differentiability, directional derivatives, subgradients and convex conjugation. We also provide results on the existence of conditional subgradients and give a conditional version of the Fenchel–Moreau theorem. In Section 8 we study conditional inf-convolutions.

Notation. We assume $\mu(\Omega) > 0$ and define $\mathcal{F}_+ := \{A \in \mathcal{F} : \mu[A] > 0\}$. By L we denote the set of all measurable functions $X : \Omega \rightarrow \mathbb{R} \cup \{\pm\infty\}$, where two of them are identified if they agree a.e. (almost everywhere). In particular, for $X, Y \in L$, $X = Y$, $X > Y$ and $X \geq Y$ will be understood in the a.e. sense. Analogously, for sets $A, B \in \mathcal{F}$, we write $A = B$ if $\mu[A \Delta B] = 0$ and $A \subseteq B$ if $\mu[A \setminus B] = 0$. The set $L^0 := \{X \in L : |X| < \infty\}$ with the a.e. order is a lattice ordered ring, and every non-empty subset C of L has a least upper bound and a greatest lower bound in L with respect to the a.e. order. We follow the usual convention in measure theory and denote them by $\text{ess sup } C$ and $\text{ess inf } C$, respectively. It is well-known (see for instance, Neveu, 1975) that there exist sequences (X_n) and (Y_n) in C such that $\text{ess sup } C = \sup_n X_n$ and $\text{ess inf } C = \inf_n Y_n$. Moreover, if C is directed upwards, (X_n) can be chosen such that $X_{n+1} \geq X_n$, and if C is directed downwards, (Y_n) can be chosen so that $Y_{n+1} \leq Y_n$. For a set $A \in \mathcal{F}$, we denote by 1_A the characteristic function of A , that is, the function $1_A : \Omega \rightarrow \{0, 1\}$ which is 1 on A and 0 elsewhere. If \mathcal{A} is a subset of \mathcal{F} , we set $\text{ess sup } \mathcal{A} := \{\text{ess sup}_{A \in \mathcal{A}} 1_A = 1\} \in \mathcal{F}$ and $\text{ess inf } \mathcal{A} := \{\text{ess inf}_{A \in \mathcal{A}} 1_A = 1\} \in \mathcal{F}$. Furthermore, we use the notation $L_+^0 := \{X \in L^0 : X \geq 0\}$, $L_{++}^0 := \{X \in L^0 : X > 0\}$, $\bar{L} := \{X \in L : X > -\infty\}$, $\underline{L} := \{X \in L : X < +\infty\}$ and $\mathbb{N} := \{1, 2, \dots\}$. By $\mathbb{N}(\mathcal{F})$ we denote the set of all measurable functions $N : \Omega \rightarrow \mathbb{N}$.

2 Algebraic structures and generating sets

We fix $d \in \mathbb{N}$ and consider the set $(L^0)^d := \{(X^1, \dots, X^d) : X^i \in L^0\}$. On $(L^0)^d$ we define the conditional inner product and conditional 2-norm by

$$\langle X, Y \rangle := \sum_{i=1}^d X^i Y^i \quad \text{and} \quad \|X\| := \langle X, X \rangle^{1/2}.$$

For every $A \in \mathcal{F}$, $1_A L^0$ is a subring of L^0 , and provided that $\mu[A] > 0$, $1_A(L^0)^d$ is a free $1_A L^0$ -module of rank d generated by the base $1_A e_i$, $i = 1, \dots, d$, where e_i is the i -th unit vector in $\mathbb{R}^d \subseteq (L^0)^d$. In particular, $(L^0)^d$ is a free L^0 -module of rank d .

Definition 2.1. We call a subset C of $(L^0)^d$

- stable if $1_A X + 1_{A^c} Y \in C$ for all $X, Y \in C$ and $A \in \mathcal{F}$;
- σ -stable if $\sum_{n \in \mathbb{N}} 1_{A_n} X_n \in C$ for every sequence $(X_n)_{n \in \mathbb{N}}$ in C and pairwise disjoint sets $A_n \in \mathcal{F}$ satisfying $\Omega = \bigcup_{n \in \mathbb{N}} A_n$;
- L^0 -convex if $\lambda X + (1 - \lambda)Y \in C$ for all $X, Y \in C$ and $\lambda \in L^0$ such that $0 \leq \lambda \leq 1$;
- an L^0 -convex cone if it is L^0 -convex and $\lambda X \in C$ for all $X \in C$ and $\lambda \in L^0_{++}$;
- L^0 -affine if $\lambda X + (1 - \lambda)Y \in C$ for all $X, Y \in C$ and $\lambda \in L^0$;
- L^0 -linear (or an L^0 -submodule) if $\lambda X + Y \in C$ for all $X, Y \in C$ and $\lambda \in L^0$.

For an arbitrary subset C of $(L^0)^d$ and $A \in \mathcal{F}$, we denote by $\text{st}_A(C)$, $\text{sst}_A(C)$, $\text{conv}_A(C)$, $\text{ccone}_A(C)$, $\text{aff}_A(C)$, $\text{lin}_A(C)$ the smallest subset of $1_A(L^0)^d$ containing $1_A C$ that is stable, σ -stable, L^0 -convex, an L^0 -convex cone, L^0 -affine, or L^0 -linear, respectively. If $A = \Omega$, we just write $\text{st}(C)$, $\text{sst}(C)$, $\text{conv}(C)$, $\text{ccone}(C)$, $\text{aff}(C)$, $\text{lin}(C)$ for these sets.

Remark 2.2. It can easily be checked that if C is a non-empty subset of $(L^0)^d$ and $A \in \mathcal{F}$, then

$$\begin{aligned} \text{st}_A(C) &= \left\{ \sum_{n=1}^k 1_{A_n} X_n : k \in \mathbb{N}, X_n \in C, A_n \in \mathcal{F}, \bigcup_{n=1}^k A_n = A, A_m \cap A_n = \emptyset \text{ for } m \neq n \right\}; \\ \text{sst}_A(C) &= \left\{ \sum_{n \in \mathbb{N}} 1_{A_n} X_n : X_n \in C, A_n \in \mathcal{F}, \bigcup_{n \in \mathbb{N}} A_n = A, A_m \cap A_n = \emptyset \text{ for } m \neq n \right\}; \\ \text{conv}_A(C) &= \left\{ \sum_{n=1}^k \lambda_n X_n : k \in \mathbb{N}, X_n \in C, \lambda_n \in 1_A L^0_+, \sum_{n=1}^k \lambda_n = 1_A \right\}; \\ \text{ccone}_A(C) &= \left\{ \sum_{n=1}^k \lambda_n X_n : k \in \mathbb{N}, X_n \in C, \lambda_n \in 1_A L^0_+, \sum_{n=1}^k \lambda_n \in 1_A L^0_{++} \right\}; \\ \text{aff}_A(C) &= \left\{ \sum_{n=1}^k \lambda_n X_n : k \in \mathbb{N}, X_n \in C, \lambda_n \in 1_A L^0, \sum_{n=1}^k \lambda_n = 1_A \right\}; \\ \text{lin}_A(C) &= \left\{ \sum_{n=1}^k \lambda_n X_n : k \in \mathbb{N}, X_n \in C, \lambda_n \in 1_A L^0 \right\}. \end{aligned}$$

It follows that if $C = \{X_1, \dots, X_k\}$ for finitely many $X_1, \dots, X_k \in (L^0)^d$, then the sets $\text{conv}_A(C)$, $\text{ccone}_A(C)$, $\text{aff}_A(C)$, $\text{lin}_A(C)$ are all σ -stable.

Definition 2.3. Let $A \in \mathcal{F}_+$ and $k \in \mathbb{N}$. We call $X_1, \dots, X_k \in (L^0)^d$ linearly independent on A if $1_A X_1, \dots, 1_A X_k$ are linearly independent in the $1_A L^0$ -module $1_A(L^0)^d$, that is, $(0, \dots, 0)$ is the only vector $(\lambda_1, \dots, \lambda_k) \in 1_A(L^0)^k$ satisfying

$$\lambda_1 X_1 + \dots + \lambda_k X_k = 0.$$

We say that X_1, \dots, X_k are orthogonal on A if $1_A \langle X_i, X_j \rangle = 0$ for $i \neq j$ and orthonormal on A if in addition, $1_A \|X_i\| = 1_A$, $1 \leq i \leq k$. If X_1, \dots, X_k are linearly independent on A and $\text{lin}_A \{X_1, \dots, X_k\} = 1_A C$ for some subset C of $(L^0)^d$, we call them a basis of C on A . If in addition, X_1, \dots, X_k are orthogonal or orthonormal on A , we say X_1, \dots, X_k is an orthogonal or orthonormal basis of C on A , respectively.

Lemma 2.4. Let $A \in \mathcal{F}$ and $X_1, \dots, X_k, Y \in (L^0)^d$ for some $k \in \mathbb{N}$. Then there exists a largest subset $B \in \mathcal{F}$ of A such that $1_B Y \in \text{lin}_B \{X_1, \dots, X_k\}$.

Proof. The set

$$\mathcal{A} := \{B \in \mathcal{F} : B \subseteq A \text{ and } 1_B Y \in \text{lin}_B \{X_1, \dots, X_k\}\}$$

is directed upwards. So it contains an increasing sequence $(B_n)_{n \in \mathbb{N}}$ such that $B := \bigcup_n B_n = \text{ess sup } \mathcal{A}$. B is the largest element of \mathcal{A} . \square

Proposition 2.5. *Let $A \in \mathcal{F}_+$ and $k, l \in \mathbb{N}$. Assume $X_1, \dots, X_k \in (L^0)^d$ are linearly independent on A and $\text{lin}_A \{X_1, \dots, X_k\} \subseteq \text{lin}_A \{Y_1, \dots, Y_l\}$ for some $Y_1, \dots, Y_l \in (L^0)^d$. Then $k \leq l$. Moreover, if $k = l$, then Y_1, \dots, Y_l are linearly independent on A and $\text{lin}_A \{X_1, \dots, X_k\} = \text{lin}_A \{Y_1, \dots, Y_l\}$.*

Proof. One can write $1_A X_1 = \sum_{i=1}^l \lambda_i 1_A Y_i$ for some $\lambda_i \in L^0$. So there exists a $\sigma(1) \in \{1, \dots, l\}$ such that $A_1 := A \cap \{\lambda_{\sigma(1)} \neq 0\} \in \mathcal{F}_+$, and one obtains

$$\text{lin}_{A_1} \{X_1, \dots, X_k\} \subseteq \text{lin}_{A_1} \{Y_1, \dots, Y_l\} = \text{lin}_{A_1} (\{X_1, Y_1, \dots, Y_l\} \setminus \{Y_{\sigma(1)}\}).$$

In particular, if $k \geq 2$, one must have $l \geq 2$, and it follows inductively that there exist $A_2, \dots, A_d \in \mathcal{F}_+$ and an injection $\sigma : \{1, \dots, k\} \rightarrow \{1, \dots, l\}$ such that for all $i \in \{1, \dots, k\}$,

$$\text{lin}_{A_i} \{X_1, \dots, X_k\} \subseteq \text{lin}_{A_i} \{Y_1, \dots, Y_l\} = \text{lin}_{A_i} (\{X_1, \dots, X_i, Y_1, \dots, Y_l\} \setminus \{Y_{\sigma(i)}, \dots, Y_{\sigma(i)}\}).$$

This shows that $k \leq l$.

Now assume $k = l$ and Y_1, \dots, Y_l are not linearly independent on A . Then there exist $B \in \mathcal{F}_+$ and $j \in \{1, \dots, k\}$ such that

$$\text{lin}_B \{X_1, \dots, X_k\} \subseteq \text{lin}_B \{Y_1, \dots, Y_k\} = \text{lin}_B (\{Y_1, \dots, Y_k\} \setminus \{Y_j\}),$$

a contradiction to the first part of the proposition. So if $k = l$, Y_1, \dots, Y_k must be linearly independent on A , and it remains to show that $\text{lin}_A \{X_1, \dots, X_k\} = \text{lin}_A \{Y_1, \dots, Y_k\}$. To do this, we assume that $\text{lin}_A \{X_1, \dots, X_k\} \subsetneq \text{lin}_A \{Y_1, \dots, Y_k\}$. Then $Y_j \notin \text{lin}_A \{X_1, \dots, X_k\}$ for at least one $j \in \{1, \dots, k\}$. By Lemma 2.4, there exists a largest subset $B \in \mathcal{F}$ of A such that $1_B Y_j \in \text{lin}_B \{X_1, \dots, X_k\}$. The set $D := A \setminus B$ is in \mathcal{F}_+ , and X_1, \dots, X_k, Y_j are linearly independent on D . But then

$$\text{lin}_D \{X_1, \dots, X_k, Y_j\} \subseteq \text{lin}_D \{Y_1, \dots, Y_k\},$$

again contradicts the first part of the proposition, and the proof is complete. \square

Corollary 2.6. *Let $A \in \mathcal{F}_+$ and $k, l \in \mathbb{N}$. Assume $X_1, \dots, X_k \in (L^0)^d$ are linearly independent on A and $\text{lin}_A \{X_1, \dots, X_k\} = \text{lin}_A \{Y_1, \dots, Y_l\}$ for some $Y_1, \dots, Y_l \in (L^0)^d$ that are also linearly independent on A . Then $k = l \leq d$, and if $k = l = d$, one has $\text{lin}_A \{X_1, \dots, X_k\} = \text{lin}_A \{Y_1, \dots, Y_l\} = 1_A (L^0)^d$.*

Proof. The corollary follows from Proposition 2.5 by noticing that

$$\text{lin}_A \{X_1, \dots, X_k\} = \text{lin}_A \{Y_1, \dots, Y_l\} \subseteq \text{lin}_A (e_1, \dots, e_d) = 1_A (L^0)^d.$$

\square

Lemma 2.7. *Let C be a non-empty σ -stable subset of $(L^0)^d$ and $X_1, \dots, X_k \in (L^0)^d$ for some $k \in \mathbb{N}$. Then for given $A \in \mathcal{F}_+$, each of the collections*

$$\{B \in \mathcal{F}_+ : B \subseteq A \text{ and there exists a } Y \in C \text{ such that } \|Y\| > 0 \text{ on } B\} \quad (2.1)$$

and

$$\{B \in \mathcal{F}_+ : B \subseteq A \text{ and there exists } Y \in C \text{ such that } X_1, \dots, X_k, Y \text{ are linearly independent on } B\} \quad (2.2)$$

is either empty or contains a largest set.

Proof. Let us denote the collection (2.1) by \mathcal{A}_1 and (2.2) by \mathcal{A}_2 . Both are directed upwards. So if either one of them is non-empty, it contains an increasing sequence of sets B_n with corresponding $Y_n \in C$, $n \in \mathbb{N}$, such that $B := \bigcup_n B_n = \text{ess sup } \mathcal{A}_i$. Since C is σ -stable,

$$Y := Y_1 1_{B_1 \cup B^c} + \sum_{n \geq 2} 1_{B_n \setminus B_{n-1}} Y_n$$

belongs to C . In the first case one has $\|Y\| > 0$ on B , and in the second one, X_1, \dots, X_k, Y are linearly independent on B . This proves the lemma. \square

Theorem 2.8. *Let C be a σ -stable subset of $(L^0)^d$ containing an element $X \neq 0$. Then there exist a unique number $k \in \{1, \dots, d\}$, unique pairwise disjoint sets $A_0, \dots, A_k \in \mathcal{F}$ and $X_1, \dots, X_k \in C$ such that the following hold:*

(i) $\bigcup_{i=0}^k A_i = \Omega$ and $\mu[A_k] > 0$;

(ii) $1_{A_0}C = \{0\}$;

(iii) *For all $i \in \{1, \dots, k\}$ satisfying $\mu[A_i] > 0$, X_1, \dots, X_i is a basis of $\text{lin}(C)$ on A_i .*

Proof. That k and the sets A_0, \dots, A_k are unique follows from Corollary 2.6. To show the existence of A_i and X_i satisfying (i)–(iii), we construct them inductively. Since C contains an element $X \neq 0$, it follows from Lemma 2.7 that there exists a largest set $B_1 \in \mathcal{F}_+$ such that $\|Y\| > 0$ on B_1 for some $Y \in C$. Choose such a Y and call it X_1 . One must have $1_{B_1^c}C = \{0\}$. If there exist no $B \in \mathcal{F}_+$ and $Y \in C$ such that X_1, Y are linearly independent on B , one obtains from Lemma 2.4 that $1_{B_1}Y \in \text{lin}_{B_1}\{X_1\}$ for all $Y \in C$, and therefore, $\text{lin}_{B_1}(C) = \text{lin}_{B_1}\{X_1\}$. So one can set $k = 1$, $A_0 = B_1^c$ and $A_1 = B_1$. On the other hand, if there exists a $B \in \mathcal{F}_+$ and $Y \in C$ such that X_1, Y are linearly independent on B , Lemma 2.7 yields a largest such set B_2 with a corresponding $X_2 \in C$. If there exists no $B \in \mathcal{F}_+$ and $Y \in C$ such that X_1, X_2, Y are linearly independent on B , then $\text{lin}_{B_2}(C) = \text{lin}_{B_2}\{X_1, X_2\}$ and one can set $k = 2$, $A_0 = B_2^c$, $A_1 = B_1 \setminus B_2$ and $A_2 = B_2$. Otherwise, one continues like this until there is no $B \in \mathcal{F}_+$ and $Y \in C$ such that X_1, \dots, X_k, Y are linearly independent on B . Such a k must exist and $k \leq d$. Otherwise one would have $X_1, \dots, X_{d+1} \in C$ that are linearly independent on some $B \in \mathcal{F}_+$, a contradiction to Corollary 2.6. One sets $A_0 = B_k^c$, $A_1 = B_1 \setminus B_2, \dots, A_{k-1} = B_{k-1} \setminus B_k$, $A_k = B_k$. \square

Corollary 2.9. *Let C be a non-empty σ -stable subset of $(L^0)^d$ and $A \in \mathcal{F}$. Then $\text{aff}_A(C)$ and $\text{lin}_A(C)$ are again σ -stable.*

Proof. If $1_A C = \{0\}$, then $\text{aff}_A(C) = \text{lin}_A(C) = \{0\}$, and the corollary is clear. Otherwise, one obtains from Theorem 2.8 that there exists a $k \in \{1, \dots, d\}$, disjoint sets $A_0, \dots, A_k \in \mathcal{F}$ and $X_1, \dots, X_k \in C$ such that $\bigcup_{i=0}^k A_i = A$, $1_{A_0}C = \{0\}$ and for all $i \in \{1, \dots, k\}$ satisfying $\mu[A_i] > 0$, X_1, \dots, X_i is a basis of $\text{lin}_A(C)$ on A_i . Now it can easily be verified that $\text{lin}_A(C)$ is σ -stable. To see that $\text{aff}_A(C)$ is σ -stable, one picks an $X \in 1_A C$. Then $\text{aff}_A(C) - X = \text{lin}_A(C - X)$ is σ -stable. So $\text{aff}_A(C)$ is σ -stable too. \square

Definition 2.10. *The orthogonal complement of a non-empty subset C of $(L^0)^d$ is given by*

$$C^\perp := \{X \in (L^0)^d : \langle X, Y \rangle = 0 \text{ for all } Y \in C\}.$$

It is clear that C^\perp is an L^0 -linear subset of $(L^0)^d$ satisfying

$$C \cap C^\perp \subseteq \{0\} \quad \text{and} \quad C \subseteq C^{\perp\perp}.$$

As a consequence of Theorem 2.8, one obtains the following corollary.

Corollary 2.11. *Let C be a non-empty σ -stable L^0 -linear subset of $(L^0)^d$. Then there exist unique pairwise disjoint sets $A_0, \dots, A_d \in \mathcal{F}$ satisfying $\bigcup_{i=0}^d A_i = \Omega$ and an orthonormal basis X_1, \dots, X_d of $(L^0)^d$ on Ω such that $1_{A_0}C = \{0\}$, $1_{A_d}C = 1_{A_d}(L^0)^d$ and*

$$1_{A_i}C = \text{lin}_{A_i}\{X_1, \dots, X_i\}, \quad 1_{A_i}C^\perp = \text{lin}_{A_i}\{X_{i+1}, \dots, X_d\} \quad \text{for } 1 \leq i \leq d-1.$$

In particular, $C + C^\perp = (L^0)^d$, $C \cap C^\perp = \{0\}$ and $C = C^{\perp\perp}$.

Proof. The uniqueness of the sets A_1, \dots, A_d follows from Corollary 2.6, and in the special case $C = \{0\}$, one can choose $A_0 = \Omega$, $A_i = \emptyset$, $X_i = e_i$, $i = 1, \dots, d$.

If C is different from $\{0\}$, it follows from Theorem 2.8 that there exist a unique number $k \in \{1, \dots, d\}$, unique pairwise disjoint sets $A_0, \dots, A_k \in \mathcal{F}$ and $Y_1, \dots, Y_k \in C$ such that $\bigcup_{i=0}^k A_i = \Omega$, $\mu[A_k] > 0$, $1_{A_0}C = \{0\}$ and for all $i \in \{1, \dots, k\}$ satisfying $\mu[A_i] > 0$, Y_1, \dots, Y_i is a basis of C on A_i . Let us define

$$U_1 := 1_{A_1 \cup \dots \cup A_k} \frac{Y_1}{\|Y_1\|} \in C$$

and

$$Z_i := Y_i - \sum_{j=1}^{i-1} \langle Y_i, U_j \rangle U_j, \quad U_i = 1_{A_i \cup \dots \cup A_k} \frac{Z_i}{\|Z_i\|} \quad \text{for } 2 \leq i \leq k.$$

Then for every $i \in \{1, \dots, k\}$ satisfying $\mu[A_i] > 0$, U_1, \dots, U_i is an orthonormal basis of C on A_i . If $k = d$, one obtains from Corollary 2.6 that $1_{A_d} C = \text{lin}_{A_d} \{U_1, \dots, U_d\} = 1_{A_d} (L^0)^d$. If $k < d$, we set $A_{k+1} = \dots = A_d = \emptyset$, and $1_{A_d} C = 1_{A_d} (L^0)^d$ holds trivially. By Corollary 2.6 and Lemma 2.7, there exist $V_i \in C$, $i = 1, \dots, d$ such that

$$1_{A_0} (L^0)^d = \text{lin}_{A_0} \{V_1, \dots, V_d\}$$

and

$$1_{A_i} (L^0)^d = \text{lin}_{A_i} \{U_1, \dots, U_i, V_{i+1}, \dots, V_d\} \quad \text{for all } i = 1, \dots, d-1.$$

Set

$$X_1 := 1_{A_1 \cup \dots \cup A_d} U_1 + 1_{A_0} \frac{V_1}{\|V_1\|}$$

and

$$W_i := V_i - \sum_{j=1}^{i-1} \langle V_i, X_j \rangle X_j, \quad X_i = 1_{A_i \cup \dots \cup A_d} U_i + 1_{A_0 \cup \dots \cup A_{i-1}} \frac{W_i}{\|W_i\|} \quad \text{for } 2 \leq i \leq d.$$

Then X_1, \dots, X_d are orthonormal on Ω such that

$$1_{A_i} C = \text{lin}_{A_i} \{X_1, \dots, X_i\}, \quad 1_{A_i} C^\perp = \text{lin}_{A_i} \{X_{i+1}, \dots, X_d\} \quad \text{for } 1 \leq i \leq d-1.$$

It is clear that $C + C^\perp = (L^0)^d$, $C \cap C^\perp = \{0\}$ and $C = C^{\perp\perp}$. □

Corollary 2.12. *Let C be a non-empty σ -stable L^0 -linear subset of $(L^0)^d$. Then every $X \in (L^0)^d$ has a unique decomposition $X = Y + Z$ for $Y \in C$, $Z \in C^\perp$, and $\|Z\| \leq \|X - V\|$ for every $V \in C$.*

Proof. That X has a unique decomposition $X = Y + Z$, $Y \in C$, $Z \in C^\perp$ is a consequence of Corollary 2.11. Moreover, if $V \in C$, then

$$\|Z\|^2 \leq \|Z\|^2 + \|Y - V\|^2 = \|Z + Y - V\|^2 = \|X - V\|^2.$$

□

3 Converging sequences, sequential closures and sequential continuity

Definition 3.1. *We call a subset C of $(L^0)^d$ sequentially closed if it contains every $X \in (L^0)^d$ that is an a.e. limit of a sequence $(X_n)_{n \in \mathbb{N}}$ in C . For an arbitrary subset C of $(L^0)^d$ and $A \in \mathcal{F}_+$, we denote by $\lim_A(C)$ the set consisting of all a.e. limits of sequences in $1_A C$ and by $\text{cl}_A(C)$ the smallest sequentially closed subset of $1_A (L^0)^d$ containing $1_A C$. In the special case $A = \Omega$, we just write $\lim(C)$ and $\text{cl}(C)$, respectively.*

Proposition 3.2. *For all subsets C of $(L^0)^d$ and $A \in \mathcal{F}_+$ one has $\lim_A(C) = \text{cl}_A(C)$.*

Proof. It is clear that $\lim_A(C) \subseteq \text{cl}_A(C)$. To show that the two sets are equal, it is enough to prove that $\lim_A(C)$ is sequentially closed. So let $(X_n)_{n \in \mathbb{N}}$ be a sequence in $\lim_A(C)$ that converges a.e. to some $X \in 1_A (L^0)^d$. Since $(\Omega, \mathcal{F}, \mu)$ is σ -finite, there exists an increasing sequence A_n , $n \in \mathbb{N}$, of measurable sets such that $\bigcup_n A_n = A$ and $\mu[A_n] < +\infty$. For every n there exists a sequence $(Y_m)_{m \in \mathbb{N}}$ in $1_{A_n} C$ converging a.e. to X_n . Therefore,

$$\mu[A_n \cap \{|Y_m - X_n| > 1/n\}] \rightarrow 0 \quad \text{for } m \rightarrow \infty,$$

and one can choose $m_n \in \mathbb{N}$ such that

$$\mu[B_n] \leq 2^{-n}, \quad \text{where } B_n = A_n \cap \{|Y_{m_n} - X_n| > 1/n\}.$$

It follows from the Borel–Cantelli lemma that $\mu \left[\bigcap_{k \in \mathbb{N}} \bigcup_{n \geq k} B_n \right] = 0$, which implies $Y_{m_n} \rightarrow X$ a.e. for $n \rightarrow \infty$. So $X \in \lim_A(C)$, and the proof is complete. □

Corollary 3.3. *If C is a stable subset of $(L^0)^d$ and $A \in \mathcal{F}_+$, then*

$$\lim_A(C) = 1_A \lim(C) = \text{cl}_A(C) = 1_A \text{cl}(C).$$

In particular, if C is stable and sequentially closed, then so is $1_A C$.

Proof. $\lim_A(C) = 1_A \lim(C)$ is a consequence of the stability of C . Moreover, it follows from Proposition 3.2 that $\lim_A(C) = \text{cl}_A(C)$ and $\lim(C) = \text{cl}(C)$. This proves the corollary. \square

Corollary 3.4. *If C is a stable subset of $(L^0)^d$ and $A \in \mathcal{F}_+$, then $\text{cl}_A(C)$ is σ -stable. Moreover, if C is L^0 -convex, an L^0 -convex cone, L^0 -affine or L^0 -linear, then so is $\text{cl}_A(C)$.*

Proof. By Proposition 3.2, $\text{cl}_A(C)$ is equal to $\lim_A(C)$. So for all $X, Y \in \text{cl}_A(C)$ there exist sequences $(X_n)_{n \in \mathbb{N}}$ and $(Y_n)_{n \in \mathbb{N}}$ in $1_A C$ such that $X_n \rightarrow X$ a.e. and $Y_n \rightarrow Y$ a.e. Since for all $B \in \mathcal{F}$, $1_B X_n + 1_{B^c} Y_n \in 1_A C$ and $1_B X_n + 1_{B^c} Y_n \rightarrow 1_B X + 1_{B^c} Y$ a.e., one obtains that $1_B X + 1_{B^c} Y$ belongs to $\lim_A(C) = \text{cl}_A(C)$. This shows that $\text{cl}_A(C)$ is stable. Since it is also sequentially closed, it must be σ -stable. The rest of the corollary follows similarly. \square

Proposition 3.5. *Every σ -stable L^0 -affine subset C of $(L^0)^d$ is sequentially closed.*

Proof. If C is empty, the corollary is trivial. Otherwise, choose $X \in C$. Then $D = C - X$ is a σ -stable L^0 -linear subset of $(L^0)^d$, and the corollary follows if we can show that D is sequentially closed. So let $(Y_n)_{n \in \mathbb{N}}$ be a sequence in D converging a.e. to some $Y \in (L^0)^d$. By Corollary 2.11, there exist unique pairwise disjoint sets $A_0, \dots, A_d \in \mathcal{F}$ satisfying $\bigcup_{i=0}^d A_i = \Omega$ and an orthonormal basis X_1, \dots, X_d of $(L^0)^d$ on Ω such that $1_{A_0} D = \{0\}$ and $1_{A_i} D = \text{lin}_{A_i} \{X_1, \dots, X_i\}$ for $1 \leq i \leq d$. Define λ_n and λ in $(L^0)^d$ by $\lambda_n^j := \langle Y_n, X_j \rangle$ and $\lambda^j := \langle Y, X_j \rangle$. Since $Y_n \rightarrow Y$ a.e., one has $\lambda_n^j \rightarrow \lambda^j$ a.e. In particular, $\lambda^j = 0$ on A_i such that $i < j$. This shows that $Y = \sum_j \lambda^j X_j \in D$. \square

The following example shows that L^0 -affine subsets of $(L^0)^d$ that are not σ -stable need not be sequentially closed.

Example 3.6. Let $\Omega = \mathbb{N}$, $\mathcal{F} = 2^{\mathbb{N}}$ and μ the counting measure. Set $X_n = 1_{\{n\}} e_1$. Then

$$\text{lin}(X_n : n \in \mathbb{N}) = \left\{ \sum_{n=1}^k \lambda_n X_n : k \in \mathbb{N}, \lambda_1, \dots, \lambda_k \in L^0 \right\}$$

is an L^0 -linear subset of $(L^0)^d$ that is not σ -stable, and $Y_k = \sum_{n=1}^k X_n$ is a sequence in $\text{lin}(X_n : n \in \mathbb{N})$ that converges a.e. to $\sum_{n \in \mathbb{N}} X_n \notin \text{lin}(X_n : n \in \mathbb{N})$. Note that $\text{lin}(X_n : n \in \mathbb{N})$ is an L^0 -submodule of $(L^0)^d$ that is not finitely generated.

The next result is a conditional version of the Bolzano–Weierstrass theorem. It is already known (see for instance, Lemma 2 in Kabanov and Stricker (2001) or Lemma 1.63 in Föllmer and Schied (2004)). But since it is important to some of our later results, we give a short proof. To state the result we need the following definition.

Definition 3.7. *We call a subset C of $(L^0)^d$ L^0 -bounded if $\text{ess sup}_{X \in C} \|X\| \in L^0$.*

Note that if $(X_n)_{n \in \mathbb{N}}$ is a sequence in $(L^0)^d$ and $N \in \mathbb{N}(\mathcal{F})$, X_N can be written as

$$X_N = \sum_{n \in \mathbb{N}} 1_{\{N=n\}} X_n.$$

In particular, X_N is in $(L^0)^d$. Moreover, if all X_n belong to a σ -stable subset C of $(L^0)^d$, then X_N is again in C .

Theorem 3.8. (Conditional version of the Bolzano–Weierstrass theorem)

Let $(X_n)_{n \in \mathbb{N}}$ be an L^0 -bounded sequence in $(L^0)^d$. Then there exists an $X \in (L^0)^d$ and a sequence $(N_n)_{n \in \mathbb{N}}$ in $\mathbb{N}(\mathcal{F})$ such that $N_{n+1} > N_n$ for all $n \in \mathbb{N}$ and $\lim_{n \rightarrow \infty} X_{N_n} = X$ a.e.

Proof. There exists a $Y \in L^0_+$ such that $\|X_n\| \leq Y$ for all $n \in \mathbb{N}$. Therefore, the a.e. limit $X^1 := \lim_{n \rightarrow \infty} \inf_{m \geq n} X_m^1$ exists and is in L^0 . Define $N_0^1 := 0$ and

$$N_n^1(\omega) := \min \{m \in \mathbb{N} : m > N_{n-1}^1(\omega) \text{ and } X_m^1(\omega) \leq X^1(\omega) + 1/n\} \in \mathbb{N}(\mathcal{F}), \quad n \in \mathbb{N}.$$

Then $N_{n+1}^1 > N_n^1$ for all $n \in \mathbb{N}$ and $\lim_{n \rightarrow \infty} X_{N_n^1}^1 = X^1$ a.e. Now set $Y_n^2 = X_{N_n^1}^2$. Then there exists a sequence $(M_n^2)_{n \in \mathbb{N}}$ in $\mathbb{N}(\mathcal{F})$ such that $M_{n+1}^2 > M_n^2$ for all $n \in \mathbb{N}$ and $\lim_{n \rightarrow \infty} Y_{M_n^2}^2 = X^2 := \lim_{n \rightarrow \infty} \inf_{m \geq n} Y_m^2$ a.e. $N_n^2 := N_{M_n^2}^1$, $n \in \mathbb{N}$, defines a sequence in $\mathbb{N}(\mathcal{F})$ satisfying $N_{n+1}^2 > N_n^2$ for all $n \in \mathbb{N}$, and one has $\lim_{n \rightarrow \infty} X_{N_n^2}^i = X^i$ a.e. for $i = 1, 2$. If one continues like this, one obtains $X^1, \dots, X^d \in L^0$ and a sequence $(N_n)_{n \in \mathbb{N}}$ in $\mathbb{N}(\mathcal{F})$ such that $N_{n+1} > N_n$ for all $n \in \mathbb{N}$ and $\lim_{n \rightarrow \infty} X_{N_n} = X = (X^1, \dots, X^d)$ a.e. \square

Corollary 3.9. *Let $(X_n)_{n \in \mathbb{N}}$ be a sequence in a sequentially closed L^0 -bounded stable subset C of $(L^0)^d$. Then there exists an $X \in C$ and a sequence $(N_n)_{n \in \mathbb{N}}$ in $\mathbb{N}(\mathcal{F})$ such that $N_{n+1} > N_n$ for all $n \in \mathbb{N}$ and $\lim_{n \rightarrow \infty} X_{N_n} = X$ a.e.*

Proof. Since $(X_n)_{n \in \mathbb{N}}$ is L^0 -bounded, it follows from Theorem 3.8 that there exists $X \in (L^0)^d$ and a sequence $(N_n)_{n \in \mathbb{N}}$ in $\mathbb{N}(\mathcal{F})$ such that $N_{n+1} > N_n$ for all $n \in \mathbb{N}$ and $\lim_{n \rightarrow \infty} X_{N_n} = X$ a.e. It remains to show that X belongs to C . By Corollary 3.4 the subset C is σ -stable. Hence, X_{N_n} belongs to C for all $n \in \mathbb{N}$, which implies that X is in C too. \square

Corollary 3.10. *Let C and D be non-empty sequentially closed stable subsets of $(L^0)^d$ such that D is L^0 -bounded. Then $C + D$ is sequentially closed and stable.*

Proof. That $C + D$ is stable is clear. To show that $C + D$ is sequentially closed, choose a sequence $(X_n)_{n \in \mathbb{N}}$ in C and a sequence $(Y_n)_{n \in \mathbb{N}}$ in D such that $X_n + Y_n \rightarrow Z$ a.e. for some $Z \in (L^0)^d$. By Theorem 3.8, there exists $Y \in D$ and a sequence $(N_n)_{n \in \mathbb{N}}$ in $\mathbb{N}(\mathcal{F})$ such that $N_{n+1} > N_n$ for all $n \in \mathbb{N}$ and $\lim_{n \rightarrow \infty} Y_{N_n} = Y$ a.e. It follows that $\lim_{n \rightarrow \infty} X_{N_n} = Z - Y$ a.e. Since C is sequentially closed, $Z - Y$ belongs to C . Hence, Z is in $C + D$. \square

Another consequence of Theorem 3.8 is that conditional Cauchy sequences converge if they are defined as follows:

Definition 3.11. *We call a sequence $(X_n)_{n \in \mathbb{N}}$ in $(L^0)^d$ L^0 -Cauchy if for every $\varepsilon \in L^0_{++}$ there exists an $N_0 \in \mathbb{N}(\mathcal{F})$ such that $\|X_{N_1} - X_{N_2}\| \leq \varepsilon$ for all $N_1, N_2 \in \mathbb{N}(\mathcal{F})$ satisfying $N_1, N_2 \geq N_0$.*

Theorem 3.12. *Every L^0 -Cauchy sequence $(X_n)_{n \in \mathbb{N}}$ in $(L^0)^d$ converges a.e. to some $X \in (L^0)^d$.*

Proof. Choose $N_0 \in \mathbb{N}(\mathcal{F})$ such that $\|X_{N_1} - X_{N_2}\| \leq 1$ for all $N_1, N_2 \in \mathbb{N}(\mathcal{F})$ satisfying $N_1, N_2 \geq N_0$. Then

$$\|X_n\| \leq 1 + \sum_{m \in \mathbb{N}} 1_{\{m \leq N_0\}} \|X_m\| \in L^0$$

for all $n \in \mathbb{N}$. So it follows from Theorem 3.8 that there exist $X \in (L^0)^d$ and a sequence $(N_n)_{n \in \mathbb{N}}$ in $\mathbb{N}(\mathcal{F})$ such that $N_{n+1} > N_n$ for all $n \in \mathbb{N}$ and $\lim_{n \rightarrow \infty} X_{N_n} = X$ a.e. But since $(X_n)_{n \in \mathbb{N}}$ is L^0 -Cauchy, one has $\lim_{n \rightarrow \infty} X_n = X$ a.e. \square

The following result gives necessary and sufficient conditions for a sequentially closed L^0 -convex subset of $(L^0)^d$ to be L^0 -bounded.

Theorem 3.13. *Let C be a sequentially closed L^0 -convex subset of $(L^0)^d$ containing 0. Then C is L^0 -bounded if and only if for any $X \in C \setminus \{0\}$ there exists a $k \in \mathbb{N}$ such that $kX \notin C$.*

Proof. Suppose that C is L^0 -bounded. Then for every $0 \neq X \in C$, there exists a $k \in \mathbb{N}$ such that $\mu[\|kX\| > \text{ess sup}_{Y \in C} \|Y\|] > 0$, and therefore $kX \notin C$.

Conversely, suppose that C is not L^0 -bounded. The sequence

$$A_n := \text{ess sup} \{B \in \mathcal{F} : \|X\| \geq n \text{ on } B \text{ for some } X \in C\}, \quad n \in \mathbb{N} \cup \{0\},$$

is decreasing with limit $A := \bigcap_n A_n$. One must have $\mu[A] > 0$, since otherwise, $\|X\| \leq \sum_{n \in \mathbb{N}} n 1_{\{A_n^c \setminus A_{n-1}^c\}} \in L^0$ for all $X \in C$. Since C is sequentially closed, L^0 -convex and therefore stable, it is σ -stable. It follows that there exists a sequence $(X_n)_{n \in \mathbb{N}}$ in C such that $\|X_n\| \geq n$ on A . Since the sequence $Y_n = 1_A X_n / \|X_n\|$

is L^0 -bounded, it follows from Theorem 3.8 that there exists $Y \in (L^0)^d$ and a sequence $(N_n)_{n \in \mathbb{N}}$ in $\mathbb{N}(\mathcal{F})$ such that $N_{n+1} > N_n$ and $\lim_{n \rightarrow \infty} Y_{N_n} = Y$ a.e. Obviously, $1_A \|Y\| = 1_A$, and in particular, $Y \neq 0$. Since C is L^0 -convex, sequentially closed and contains 0, one has for all $n \geq k$,

$$kY_{N_n} = 1_A \frac{k}{\|X_{N_n}\|} X_{N_n} \in C.$$

But $\lim_{n \rightarrow \infty} kY_{N_n} = kY$. So $kY \in C$ for all $k \in \mathbb{N}$. \square

Definition 3.14. Let C be a non-empty subset of $(L^0)^d$ and $k \in \mathbb{N}$. We call a function $f : C \rightarrow (L^0)^k$

- sequentially continuous at $X \in C$ if $f(X_n) \rightarrow f(X)$ a.e. for every sequence $(X_n)_{n \in \mathbb{N}}$ in C converging to X a.e.;
- sequentially continuous if it is sequentially continuous at every $X \in C$;
- L^0 -affine if $f(\lambda X + (1 - \lambda)Y) = \lambda f(X) + (1 - \lambda)f(Y)$ for all $X, Y \in (L^0)^d$ and $\lambda \in L^0$ such that $\lambda X + (1 - \lambda)Y \in C$;
- L^0 -linear if $f(\lambda X + Y) = \lambda f(X) + f(Y)$ for all $X, Y \in (L^0)^d$ and $\lambda \in L^0$ such that $\lambda X + Y \in C$.
- We define the conditional norm of f by $\|f\| := \text{ess sup}_{X \in C, \|X\| \leq 1} \|f(X)\| \in \bar{L}$.

Proposition 3.15. Let C be a non-empty σ -stable L^0 -linear subset of $(L^0)^d$. Then $\|f\| \in L_+^0$ for every L^0 -linear function $f : C \rightarrow (L^0)^k$, $k \in \mathbb{N}$.

Proof. By Corollary 2.11, there exist unique pairwise disjoint sets $A_0, \dots, A_d \in \mathcal{F}$ satisfying $\bigcup_{i=0}^d A_i = \Omega$ and an orthonormal basis X_1, \dots, X_d of $(L^0)^d$ on Ω such that $1_{A_0} C = \{0\}$ and $1_{A_i} C = \text{lin}_{A_i} \{X_1, \dots, X_i\}$ for $1 \leq i \leq d$. For every $X \in C$ there exists a unique $\lambda \in (L^0)^d$ such that $X = \sum_{j=1}^d \lambda_j X_j$. On the set A_0 one has $f(X) = X = 0$, and on A_i for $1 \leq i \leq d$, $\|X\| = \left(\sum_{j=1}^i \lambda_j^2\right)^{1/2}$ as well as

$$\|f(X)\| = \left\| \sum_{j=1}^i \lambda_j f(X_j) \right\| \leq \sum_{j=1}^i |\lambda_j| \|f(X_j)\| \leq \left(\sum_{j=1}^i \lambda_j^2 \right)^{1/2} \left(\sum_{j=1}^i \|f(X_j)\|^2 \right)^{1/2}.$$

Therefore, $\|f\| \leq \sum_{i=1}^d 1_{A_i} \left(\sum_{j=1}^i \|f(X_j)\|^2 \right)^{1/2}$. \square

Corollary 3.16. Let C be a non-empty σ -stable L^0 -affine subset of $(L^0)^d$. Then every L^0 -affine function $f : C \rightarrow (L^0)^k$, $k \in \mathbb{N}$, is sequentially continuous.

Proof. Choose an $X_0 \in C$. Then $D = C - X_0$ is a non-empty σ -stable L^0 -linear subset of $(L^0)^d$ and $g(X) = f(X + X_0) - f(X_0)$ is an L^0 -linear function on D . By Proposition 3.15, one has $\|g\| \in L_+^0$. Moreover, $\|f(X) - f(Y)\| = \|g(X - Y)\| \leq \|g\| \|X - Y\|$, and it follows that f is sequentially continuous. \square

Corollary 3.17. Let C be a non-empty sequentially closed subset of a non-empty σ -stable L^0 -affine subset D of $(L^0)^d$. Then for every injective L^0 -affine function $f : D \rightarrow (L^0)^k$, $k \in \mathbb{N}$, $f(C)$ is a sequentially closed subset of $(L^0)^k$.

Proof. Pick an $X_0 \in C$. The corollary follows if we can show that $f(C) - f(X_0)$ is sequentially closed. So by replacing C with $C - X_0$, D with $D - X_0$ and f with $f(X + X_0) - f(X_0)$, one can assume that $X_0 = 0$, D is a σ -stable L^0 -linear subset of $(L^0)^d$ and f is injective L^0 -linear. By Corollary 3.16, f is sequentially continuous. Therefore, $f(D)$ is a non-empty σ -stable L^0 -linear subset of $(L^0)^k$, and it follows from Proposition 3.5 that it is sequentially closed. Since $f^{-1} : f(D) \rightarrow D$ is again L^0 -linear, it is also sequentially continuous. So if $(Y_n)_{n \in \mathbb{N}}$ is a sequence in $f(C)$ converging a.e. to some $Y \in (L^0)^k$, then $Y \in f(D)$ and $f^{-1}(Y_n)$ is a sequence in C converging a.e. to $f^{-1}(Y) \in D$. It follows that $f^{-1}(Y) \in C$ and $Y = f(f^{-1}(Y)) \in f(C)$. \square

Lemma 3.18. Let C be a non-empty σ -stable L^0 -linear subset of $(L^0)^d$ and $k \in \mathbb{N}$. Then every L^0 -linear function $f : C \rightarrow (L^0)^k$ has an L^0 -linear extension $F : (L^0)^d \rightarrow (L^0)^k$ such that $\|f\| = \|F\|$.

Proof. By Corollary 2.12, every $X \in (L^0)^d$ has a unique decomposition $X = Y + Z$ such that $Y \in C$ and $Z \in C^\perp$. $F(X) := f(Y)$ defines an L^0 -linear extension of f to $(L^0)^d$ such that $\|f\| = \|F\|$. \square

4 Conditional optimization

Definition 4.1. Let C be a non-empty subset of $(L^0)^d$. We call a function $f : C \rightarrow L$

- sequentially lsc (lower semicontinuous) at $X \in C$ if $f(X) \leq \liminf_{n \rightarrow \infty} f(X_n)$ for every sequence $(X_n)_{n \in \mathbb{N}}$ in C with a.e. limit X ;
- sequentially lsc if it is sequentially lsc at every $X \in C$;
- sequentially usc (upper semicontinuous) at $X \in C$ if $-f$ is sequentially lsc at X ;
- sequentially usc if it is sequentially usc at every $X \in C$;
- sequentially continuous at $X \in C$ if it is sequentially lsc and usc at X ;
- sequentially continuous if it is sequentially continuous at every $X \in C$.

In the following definition $+\infty - \infty$ is understood as $+\infty$ and $0 \cdot (\pm\infty)$ as 0.

Definition 4.2. Let $f : C \rightarrow L$ be a function on a non-empty subset C of $(L^0)^d$.

- If C is stable, we call f stable if

$$f(1_A X + 1_{A^c} Y) = 1_A f(X) + 1_{A^c} f(Y)$$

for all $X, Y \in C$ and $A \in \mathcal{F}_+$;

- If C is L^0 -convex, we call f L^0 -convex if

$$f(\lambda X + (1 - \lambda)Y) \leq \lambda f(X) + (1 - \lambda)f(Y)$$

for all $X, Y \in C$ and $\lambda \in L^0$ such that $0 \leq \lambda \leq 1$;

- If C is L^0 -convex, we call f strictly L^0 -convex if

$$f(\lambda X + (1 - \lambda)Y) < \lambda f(X) + (1 - \lambda)f(Y) \quad \text{on the set } \{X \neq \lambda X + (1 - \lambda)Y\} \neq Y\}$$

for all $X, Y \in C$ and $\lambda \in L^0$ such that $0 \leq \lambda \leq 1$.

Lemma 4.3. Let $f : C \rightarrow L$ be an L^0 -convex function on an L^0 -convex subset C of $(L^0)^d$. Then f is also stable.

Proof. Let $X, Y \in C$ and $A \in \mathcal{F}_+$. Denote $Z = 1_A X + 1_{A^c} Y$. Then one has $1_A f(Z) \leq 1_A f(X)$ and $1_A f(X) = 1_A f(1_A Z + 1_{A^c} X) \leq 1_A f(Z)$. This shows that $1_A f(Z) = 1_A f(X)$. Analogously, one obtains $1_{A^c} f(Z) = 1_{A^c} f(Y)$ and therefore $f(Z) = 1_A f(X) + 1_{A^c} f(Y)$. \square

Theorem 4.4. Let C be a sequentially closed stable subset of $(L^0)^d$ and $f : C \rightarrow \bar{L}$ a sequentially lsc stable function. Assume there exists an $X_0 \in C$ such that the set

$$\{X \in C : f(X) \leq f(X_0)\}$$

is L^0 -bounded. Then there exists an $\hat{X} \in C$ such that

$$f(\hat{X}) = \operatorname{ess\,inf}_{X \in C} f(X).$$

If C and f are L^0 -convex, then the set

$$\{X \in C : f(X) = f(\hat{X})\}$$

is L^0 -convex. If in addition, f is strictly L^0 -convex, then

$$\{X \in C : f(X) = f(\hat{X})\} = \{\hat{X}\}.$$

Proof. The set $D := \{X \in C : f(X) \leq f(X_0)\}$ is sequentially closed, stable and L^0 -bounded. It follows that $\{f(X) : X \in D\}$ is directed downwards. Therefore, there exists a sequence $(X_n)_{n \in \mathbb{N}}$ in D such that $f(X_n)$ decreases a.e. to $I := \text{ess inf}_{X \in D} f(X)$. By Corollary 3.9, there exists a sequence $(N_n)_{n \in \mathbb{N}}$ in $\mathbb{N}(\mathcal{F})$ such that $N_{n+1} > N_n$ for all $n \in \mathbb{N}$ and $\lim_{n \rightarrow \infty} X_{N_n} = \hat{X}$ a.e. for some $\hat{X} \in D$. Since X_{N_n} belongs to D and

$$f(X_{N_n}) = \sum_{m \geq n} 1_{\{N_n=m\}} f(X_m) \leq f(X_n) \quad \text{for all } n,$$

one obtains from the L^0 -lower semicontinuity of f that

$$f(\hat{X}) \leq \liminf_{n \rightarrow \infty} f(X_{N_n}) \leq \lim_{n \rightarrow \infty} f(X_n) = I.$$

This shows the first part of the theorem. That $\{X \in C : f(X) = f(\hat{X})\}$ is L^0 -convex if C and f are L^0 -convex, is clear. Finally, assume C is L^0 -convex and f strictly L^0 -convex. Then if there exists an X in C such that $f(X) = f(\hat{X})$, one has

$$f\left(\frac{X + \hat{X}}{2}\right) < \frac{f(X) + f(\hat{X})}{2}$$

on the set $\{X \neq \hat{X}\}$. It follows that $\mu[X \neq \hat{X}] = 0$. \square

Corollary 4.5. *Let C and D be non-empty sequentially closed stable subsets of $L^0(\mathcal{F})^d$ such that D is L^0 -bounded. Then there exist $\hat{X} \in C$ and $\hat{Y} \in D$ such that*

$$\|\hat{X} - \hat{Y}\| = \text{ess inf}_{X \in C, Y \in D} \|X - Y\|. \quad (4.1)$$

If in addition, C and D are L^0 -convex, then $\hat{X} - \hat{Y}$ is unique.

Proof. By Corollary 3.10, the set $E = C - D$ is sequentially closed and stable. Moreover, $Z \mapsto \|Z\|$ is a sequentially continuous L^0 -convex function from E to L^0 , and for every $Z_0 \in E$, the set $\{Z \in E : \|Z\| \leq \|Z_0\|\}$ is L^0 -bounded. So one obtains from Theorem 4.4 that there exists a $\hat{Z} \in E$ such that $\|\hat{Z}\| = \text{ess inf}_{Z \in E} \|Z\|$. This shows that there exist $\hat{X} \in C$ and $\hat{Y} \in D$ satisfying (4.1). If C and D are L^0 -convex, then so is E , and for every $Z \in E$ satisfying $\|Z\| = \|\hat{Z}\|$, one has $(Z + \hat{Z})/2 \in E$ and $\|(Z + \hat{Z})/2\| < \|\hat{Z}\|$ on the set $\{Z \neq \hat{Z}\}$. It follows that $\mu[Z \neq \hat{Z}] = 0$, and the proof is complete. \square

5 Interior, relative interior and L^0 -open sets

Definition 5.1. *Let C be a non-empty subset of $(L^0)^d$ and $A \in \mathcal{F}_+$.*

- For $X \in (L^0)^d$ and $\varepsilon \in L^0_{++}$, we denote

$$B_A^\varepsilon(X) := \{Y \in 1_A(L^0)^d : 1_A \|Y - X\| \leq \varepsilon\}.$$

- The interior $\text{int}_A(C)$ of C on A consists of elements $X \in 1_A C$ for which there exists an $\varepsilon \in L^0_{++}$ such that $B_A^\varepsilon(X) \subseteq 1_A C$. If $A = \Omega$, we just write $\text{int}(C)$ for $\text{int}_A(C)$.
- The relative interior $\text{ri}_A(C)$ of C on A consists of elements $X \in 1_A C$ for which there exists an $\varepsilon \in L^0_{++}$ such that $B_A^\varepsilon(X) \cap \text{aff}_A(C) \subseteq 1_A(C)$. If $A = \Omega$, we write $\text{ri}(C)$ instead of $\text{ri}_A(C)$.
- We say C is L^0 -open on A if $1_A C = \text{int}_A(C)$. We call it L^0 -open if it is L^0 -open on Ω .

Note that one always has $1_A \text{int}(C) \subseteq \text{int}_A(C)$ but not necessarily the other way around. The collection of all L^0 -open subsets of $(L^0)^d$ forms a topology. It is studied in Filipović et al. (2009) and is related to (ε, λ) -topologies on random locally convex modules (see Guo, 2010). We point out that sequentially closed sets in $(L^0)^d$ are different from complements of L^0 -open sets. But one has the following relation between the two:

Lemma 5.2. *Let C be a σ -stable subset of $(L^0)^d$. Then $\text{cl}(C) \cap \text{int}(C^c) = \emptyset$.*

Proof. Assume $X \in \text{cl}(C) \cap \text{int}(C^c)$. By Proposition 3.2, there exists a sequence $(X_n)_{n \in \mathbb{N}}$ in C such that $X_n \rightarrow X$ a.e. On the other hand, there is an $\varepsilon \in L_{++}^0$ such that $Y \in C^c$ for every $Y \in (L^0)^d$ satisfying $\|X - Y\| \leq \varepsilon$. $N(\omega) := \min \{n \in \mathbb{N} : \|X_n(\omega) - X(\omega)\| \leq \varepsilon(\omega)\}$ is an element of $\mathbb{N}(\mathcal{F})$, and since C is σ -stable, X_N belongs to C . But at the same time one has $\|X_N - X\| \leq \varepsilon$, implying $X_N \in C^c$. This yields a contradiction. So $\text{cl}(C) \cap \text{int}(C^c) = \emptyset$. \square

Lemma 5.3. *Let C be a non-empty L^0 -convex subset of $(L^0)^d$, $A \in \mathcal{F}_+$ and $\lambda \in L^0$ such that $0 < \lambda \leq 1$. Then*

$$\lambda X + (1 - \lambda)Y \in \text{int}_A(C) \quad \text{for all } X \in \text{int}_A(C), Y \in 1_A C \quad (5.1)$$

and

$$\lambda X + (1 - \lambda)Y \in \text{ri}_A(C) \quad \text{for all } X \in \text{ri}_A(C), Y \in 1_A C. \quad (5.2)$$

If in addition, C is σ -stable, then (5.1) and (5.2) also hold for $Y \in \text{cl}_A(C)$.

Proof. Let $X \in \text{int}_A(C)$ and $Y \in 1_A C$. There exists an $\varepsilon \in L_{++}^0$ such that $B_A^\varepsilon(X)$ is contained in $1_A C$. So

$$\lambda X + (1 - \lambda)Y + Z = \lambda(X + Z/\lambda) + (1 - \lambda)Y \subseteq 1_A C$$

for all $Z \in B_A^{\varepsilon\lambda}(0)$. This shows (5.1).

To prove (5.2), we assume that $X \in \text{ri}_A(C)$ and $Y \in 1_A C$. There exists an $\varepsilon \in L_{++}^0$ such that $B_A^\varepsilon(X) \cap \text{aff}_A(C) \subseteq 1_A C$. Choose $Z \in B_A^{\varepsilon\lambda}(0)$ such that

$$\lambda X + (1 - \lambda)Y + Z \in \text{aff}_A(C).$$

Then $X + Z/\lambda \in \text{aff}_A(C)$, and therefore $X + Z/\lambda \in 1_A C$. It follows that

$$\lambda X + (1 - \lambda)Y + Z = \lambda(X + Z/\lambda) + (1 - \lambda)Y \subseteq 1_A C.$$

This shows (5.2).

If C is σ -stable, $X \in \text{int}_A(C)$ and $Y \in \text{cl}_A(C)$, there exists an $\varepsilon \in L_{++}^0$ such that $B_A^{2\varepsilon}(X) \subseteq 1_A C$. From Lemma 3.2 we know that there exists a sequence $(Y_n)_{n \in \mathbb{N}}$ in $1_A C$ converging a.e. to Y . $N(\omega) := \min \{n \in \mathbb{N} : (1 - \lambda(\omega))\|Y(\omega) - Y_n(\omega)\| \leq \lambda(\omega)\varepsilon(\omega)\}$ belongs to $\mathbb{N}(\mathcal{F})$, and Y_N is an element of C satisfying $(1 - \lambda)\|Y - Y_N\| \leq \lambda\varepsilon$. So for $Z \in B_A^{\lambda\varepsilon}(0)$, one has

$$\lambda X + (1 - \lambda)Y + Z = \lambda \left(X + \frac{(1 - \lambda)}{\lambda}(Y - Y_N) + \frac{1}{\lambda}Z \right) + (1 - \lambda)Y_N \in 1_A C,$$

which shows that $\lambda X + (1 - \lambda)Y \in \text{int}_A(C)$.

If X is in $\text{ri}_A(C)$ instead of $\text{int}_A(C)$, there exists an $\varepsilon \in L_{++}^0$ such that $B_A^{2\varepsilon}(X) \cap \text{aff}_A(C) \subseteq 1_A C$. Let $Z \in B_A^{\lambda\varepsilon}(0)$ such that

$$\lambda X + (1 - \lambda)Y + Z \in \text{aff}_A(C),$$

then

$$X + \frac{(1 - \lambda)}{\lambda}(Y - Y_N) + \frac{1}{\lambda}Z \in \text{aff}_A(C).$$

Hence

$$X + \frac{(1 - \lambda)}{\lambda}(Y - Y_N) + \frac{1}{\lambda}Z \in 1_A C,$$

and it follows that

$$\lambda X + (1 - \lambda)Y + Z = \lambda \left(X + \frac{(1 - \lambda)}{\lambda}(Y - Y_N) + \frac{1}{\lambda}Z \right) + (1 - \lambda)Y_N \in 1_A C.$$

So $\lambda X + (1 - \lambda)Y \in \text{ri}_A(C)$, and the proof is complete. \square

Corollary 5.4. *Let C be an L^0 -convex subset of $(L^0)^d$ and $A \in \mathcal{F}_+$. Then $\text{int}_A(C)$ and $\text{ri}_A(C)$ are again L^0 -convex.*

Proof. Since C is stable, it follows from Lemma 5.3 that for $X, Y \in \text{int}_A(C)$ and $\lambda \in L^0$ satisfying $0 \leq \lambda \leq 1$, one has

$$\lambda X + (1 - \lambda)Y = 1_{\{\lambda > 0\}}(\lambda X + (1 - \lambda)Y) + 1_{\{\lambda = 0\}}Y \in \text{int}_A(C).$$

This shows that $\text{int}_A(C)$ is L^0 -convex. The same argument shows that $\text{ri}_A(C)$ is L^0 -convex. \square

Definition 5.5. Let $A \in \mathcal{F}_+$. We call a subset C of $(L^0)^d$

- an L^0 -hyperplane on A if $1_A C = \{X \in 1_A(L^0)^d : \langle X, Z \rangle = V\}$
- an L^0 -halfspace on A if $1_A C = \{X \in 1_A(L^0)^d : \langle X, Z \rangle \geq V\}$

for some $V \in 1_A L^0$ and $Z \in 1_A(L^0)^d$ such that $\|Z\| > 0$ on A .

Lemma 5.6. A subset C of $(L^0)^d$ is an L^0 -hyperplane on $A \in \mathcal{F}_+$ if and only if there exist $X_0 \in 1_A(L^0)^d$ and an orthonormal basis X_1, \dots, X_d of $(L^0)^d$ on A such that

$$1_A C = \left\{ X_0 + \sum_{i=1}^{d-1} \lambda_i X_i : \lambda_i \in 1_A L^0 \right\}. \quad (5.3)$$

Similarly, C is an L^0 -halfspace on $A \in \mathcal{F}_+$ if and only if there exist $X_0 \in 1_A(L^0)^d$ and an orthonormal basis X_1, \dots, X_d of $(L^0)^d$ on A such that

$$1_A C = \left\{ X_0 + \sum_{i=1}^d \lambda_i X_i : \lambda_i \in 1_A L^0, \lambda_d \geq 0 \right\}. \quad (5.4)$$

Proof. If $1_A C$ is of the form (5.3), then $1_A C = \{X \in 1_A(L^0)^d : \langle X, X_d \rangle = \langle X_0, X_d \rangle\}$. Now assume that $1_A C = \{X \in 1_A(L^0)^d : \langle X, Z \rangle = V\}$ for some $V \in 1_A L^0$ and $Z \in 1_A(L^0)^d$ such that $\|Z\| > 0$ on A . By Corollary 2.11, there exists an orthonormal basis X_1, \dots, X_d of $(L^0)^d$ on A such that $1_A Z^\perp = \text{lin}_A \{X_1, \dots, X_{d-1}\}$ and $X_d = 1_A Z / \|Z\|$. Choose $X_0 \in 1_A(L^0)^d$ such that $\langle X_0, Z \rangle = V$. Then $1_A C$ is of the form (5.3). That C is an L^0 -halfspace on $A \in \mathcal{F}_+$ if and only if $1_A C$ is of the form (5.4) follows similarly. \square

Lemma 5.7. Let C be a σ -stable L^0 -convex subset of $(L^0)^d$ and $A \in \mathcal{F}_+$. Then $\text{int}_A(C) \neq \emptyset$ if and only if $\text{aff}_A(C) = 1_A(L^0)^d$.

Proof. Let us first assume that $X_0 \in \text{int}_A(C)$. Then $0 \in \text{int}_A(C - X_0)$, and it follows that

$$\text{aff}_A(C) = \text{aff}_A(C - X_0) + X_0 = \text{lin}_A(C - X_0) + X_0 = 1_A(L^0)^d + X_0 = 1_A(L^0)^d.$$

On the other hand, if $\text{aff}_A(C) = 1_A(L^0)^d$, choose $X_0 \in 1_A C$. Then

$$\text{lin}_A(C - X_0) = \text{aff}_A(C - X_0) = \text{aff}_A(C) - X_0 = 1_A(L^0)^d.$$

So it follows from Theorem 2.8 that there exist X_1, \dots, X_d in $1_A C$ such that $X_i - X_0$, $i = 1, \dots, d$, form a basis of $(L^0)^d$ on A . Set

$$\hat{X} := \frac{1}{d+1} \sum_{i=0}^d X_i.$$

It follows from Corollary 2.11 and Lemma 5.6 that for every $i = 0, \dots, d$, there exist $V_i \in L^0$ and $Z_i \in (L^0)^d$ such that for all $j \neq i$,

$$\langle \hat{X}, Z_i \rangle > V_i = \langle X_j, Z_i \rangle \text{ on } A.$$

This shows that $\hat{X} \in \text{int}_A \{X \in 1_A(L^0)^d : \langle X, Z_i \rangle \geq V_i\}$ for all i , which implies $\hat{X} \in \text{int}_A(C)$ since

$$\bigcap_{i=0}^d \{X \in 1_A(L^0)^d : \langle X, Z_i \rangle \geq V_i\} = \text{conv}_A \{X_0, \dots, X_d\} \subseteq 1_A C.$$

\square

6 Separation by L^0 -hyperplanes

In this section we prove results on the separation of two L^0 -convex sets in $(L^0)^d$ by an L^0 -hyperplane. As a corollary we obtain a version of the Hahn–Banach extension theorem. Hahn–Banach extension and separation results have been proved in more general modules; see e.g., Filipović et al. (2009), Guo (2010) and the references therein. However, due to the special form of $(L^0)^d$, we here are able to derive analogs of results that hold in \mathbb{R}^d but not in infinite-dimensional vector spaces. Moreover, we do not need Zorn’s lemma or the axiom of choice.

Theorem 6.1. (Strong separation)

Let C and D be non-empty L^0 -convex subsets of $(L^0)^d$. Then there exists $Z \in (L^0)^d$ such that

$$\operatorname{ess\,inf}_{X \in C} \langle X, Z \rangle > \operatorname{ess\,sup}_{Y \in D} \langle Y, Z \rangle \quad (6.1)$$

if and only if $0 \notin \operatorname{cl}_A(C - D)$ for all $A \in \mathcal{F}_+$.

Proof. Let us first assume that there exists an $A \in \mathcal{F}_+$ such that $0 \in \operatorname{cl}_A(C - D)$. From Proposition 3.2 we know that $\operatorname{cl}_A(C - D) = \lim_A(C - D)$. So there exists a sequence $(X_n)_{n \in \mathbb{N}}$ in $1_A(C - D)$ such that $X_n \rightarrow 0$ a.e. It follows that there can exist no $Z \in (L^0)^d$ satisfying (6.1).

Now assume $0 \notin \operatorname{cl}_A(C - D)$ for all $A \in \mathcal{F}_+$. It follows from Corollary 3.4 that $\operatorname{cl}(C - D)$ is L^0 -convex. So one obtains from Corollary 4.5 that there exists a $Z \in \operatorname{cl}(C - D)$ such that

$$\|Z\|^2 \leq \|(1 - \lambda)Z + \lambda W\|^2 = \|Z\|^2 + 2\lambda \langle Z, W - Z \rangle + \lambda^2 \|W - Z\|^2$$

for all $W \in \operatorname{cl}(C - D)$ and $\lambda \in L^0$ such that $0 < \lambda \leq 1$. Division by 2λ and sending λ to 0 yields $\langle W, Z \rangle \geq \|Z\|^2$. In particular,

$$\langle W, Z \rangle \geq \|Z\|^2 \quad \text{for all } W \in C - D,$$

and therefore,

$$\operatorname{ess\,inf}_{X \in C} \langle X, Z \rangle \geq \operatorname{ess\,sup}_{Y \in D} \langle Y, Z \rangle + \|Z\|^2.$$

It remains to show that $\|Z\| > 0$. But if this were not the case, the set $A = \{Z = 0\}$ would belong to \mathcal{F}_+ and $1_A Z = 0$. However, by assumption and Corollary 3.3, one has $0 \notin \operatorname{cl}_A(C - D) = 1_A \operatorname{cl}(C - D)$ for all $A \in \mathcal{F}_+$, a contradiction. \square

Corollary 6.2. Let C and D be non-empty sequentially closed L^0 -convex subsets of $(L^0)^d$ such that D is L^0 -bounded and $1_A C$ is disjoint from $1_A D$ for all $A \in \mathcal{F}_+$. Then there exists a $Z \in (L^0)^d$ such that

$$\operatorname{ess\,inf}_{X \in C} \langle X, Z \rangle > \operatorname{ess\,sup}_{Y \in D} \langle Y, Z \rangle.$$

Proof. $C - D$ is a non-empty L^0 -convex set, which by Corollary 3.10 is sequentially closed. It follows from the assumptions that $0 \notin 1_A(C - D)$ for all $A \in \mathcal{F}_+$, and we know from Corollary 3.3 that $1_A(C - D) = \operatorname{cl}_A(C - D)$. So the corollary is a consequence of Theorem 6.1. \square

Lemma 6.3. Let C be a non-empty σ -stable L^0 -convex cone in $(L^0)^d$ such that $1_A C \neq 1_A(L^0)^d$ for all $A \in \mathcal{F}_+$. Then there exists a $Z \in (L^0)^d$ such that

$$\|Z\| > 0 \quad \text{and} \quad \operatorname{ess\,inf}_{X \in C} \langle X, Z \rangle \geq 0. \quad (6.2)$$

Proof. If $C = \{0\}$, the lemma is clear. Otherwise one obtains from Theorem 2.8 that there exist $A \in \mathcal{F}$ and $X_1, \dots, X_{d-1} \in C$ such that $\operatorname{lin}_A(C) = \operatorname{lin}_A(L^0)^d$ and $\operatorname{lin}_{A^c}(C) \subseteq \operatorname{lin}_{A^c}\{X_1, \dots, X_{d-1}\}$. By Corollary 2.11, there exists $W \in \operatorname{lin}_{A^c}\{X_1, \dots, X_{d-1}\}^\perp$ such that $\|W\| > 0$ on A^c . If $\mu[A] = 0$, then $Z = W$ satisfies (6.2), and the proof is complete. If $\mu[A] > 0$, one notes that since C is an L^0 -convex cone, one has $\operatorname{aff}_A(C) = \operatorname{lin}_A(C) = 1_A(L^0)^d$. It follows from Lemma 5.7 that there exists a $Y \in \operatorname{int}_A(C)$. Then $1_B Y \in \operatorname{int}_B(C)$ for every subset $B \in \mathcal{F}_+$ of A . But this implies that $-1_B Y$ cannot be in $\operatorname{cl}_B(C)$. Otherwise it would follow from Lemma 5.3 that 0 belongs to $\operatorname{int}_B(C)$, implying that $1_B C = 1_B(L^0)^d$ and contradicting the assumptions. So Theorem 6.1 applied to $1_A C$ and $\{-Y\}$ viewed as subsets of $1_A(L^0)^d$ yields a $V \in 1_A(L^0)^d$ such that

$$\operatorname{ess\,inf}_{X \in 1_A C} \langle X, V \rangle > \langle -Y, V \rangle \quad \text{on } A.$$

Since C is an L^0 -convex cone, $Z = 1_A V + 1_{A^c} W$ satisfies condition (6.2). \square

Theorem 6.4. (Weak separation)

Let C and D be non-empty σ -stable L^0 -convex subsets of $(L^0)^d$. Then there exists a $Z \in (L^0)^d$ such that

$$\|Z\| > 0 \quad \text{and} \quad \operatorname{ess\,inf}_{X \in C} \langle X, Z \rangle \geq \operatorname{ess\,sup}_{Y \in D} \langle Y, Z \rangle \quad (6.3)$$

if and only if $0 \notin \operatorname{int}_A(C - D)$ for all $A \in \mathcal{F}_+$.

Proof. If there is an $A \in \mathcal{F}_+$ such that $0 \in \operatorname{int}_A(C - D)$, there can exist no $Z \in (L^0)^d$ such that (6.3) holds. Hence, (6.3) implies $0 \notin \operatorname{int}_A(C - D)$ for all $A \in \mathcal{F}_+$.

To show the converse implication, assume that $0 \notin \operatorname{int}_A(C - D)$ for all $A \in \mathcal{F}_+$. Clearly, $C - D$ is σ -stable and L^0 -convex. Therefore, one has $\operatorname{ccone}(C - D) = \{\lambda X : \lambda \in L_{++}^0, X \in C - D\}$, from which it can be seen that $\operatorname{ccone}(C - D)$ is σ -stable and satisfies $1_A \operatorname{ccone}(C - D) \neq 1_A(L^0)^d$ for all $A \in \mathcal{F}_+$. So one obtains from Lemma 6.3 that there exists a $Z \in (L^0)^d$ such that

$$\|Z\| > 0 \quad \text{and} \quad \operatorname{ess\,inf}_{X \in E} \langle X, Z \rangle \geq 0.$$

This implies (6.3). □

Corollary 6.5. Let C and D be two non-empty σ -stable L^0 -convex subsets of $(L^0)^d$ such that $1_A C$ is disjoint from $1_A D$ for all $A \in \mathcal{F}_+$ and D is L^0 -open. Then there exists a $Z \in (L^0)^d$ such that

$$\operatorname{ess\,inf}_{X \in C} \langle X, Z \rangle > \langle Y, Z \rangle \quad \text{for all } Y \in D.$$

Proof. It follows from Theorem 6.4 that there exists a $Z \in (L^0)^d$ such that

$$\|Z\| > 0 \quad \text{and} \quad \operatorname{ess\,inf}_{X \in C} \langle X, Z \rangle \geq \operatorname{ess\,sup}_{V \in D} \langle V, Z \rangle,$$

and since D is L^0 -open, one has

$$\operatorname{ess\,sup}_{V \in D} \langle V, Z \rangle > \langle Y, Z \rangle \quad \text{for all } Y \in D. \quad \square$$

As another consequence of Theorem 6.4 we obtain a conditional version of the Hahn–Banach extension theorem.

Corollary 6.6. (Conditional version of the Hahn–Banach extension theorem)

Let $f : (L^0)^d \rightarrow L^0$ be an L^0 -convex function such that $f(\lambda X) = \lambda f(X)$ for all $\lambda \in L_+^0$ and $g : E \rightarrow L^0$ an L^0 -linear mapping on a σ -stable L^0 -linear subset E of $(L^0)^d$ such that $g(X) \leq f(X)$ for all $X \in E$. Then there exists an L^0 -linear extension $h : (L^0)^d \rightarrow L^0$ of g such that $h(X) \leq f(X)$ for all $X \in (L^0)^d$.

Proof. Note that

$$C := \{(X, V) \in (L^0)^d \times L^0 : f(X) \leq V\} \quad \text{and} \quad D := \{(Y, g(Y)) : Y \in E\}$$

are L^0 -convex sets in $(L^0)^d \times L^0$. By Lemma 4.3, f and g are stable. It follows that C and D are σ -stable. Moreover, since $C - D$ is an L^0 -convex cone and $1_A(0, -1) \notin 1_A(C - D)$ for all $A \in \mathcal{F}_+$, one has $(0, 0) \notin \operatorname{int}_A(C - D)$ for all $A \in \mathcal{F}_+$. So one obtains from Theorem 6.4 that there exists a pair $(Z, W) \in (L^0)^d \times L^0$ such that

$$\|Z\| + |W| > 0 \quad \text{and} \quad \operatorname{ess\,inf}_{(X, V) \in C} \{\langle X, Z \rangle + VW\} \geq \operatorname{ess\,sup}_{Y \in E} \{\langle Y, Z \rangle + g(Y)W\}. \quad (6.4)$$

It follows that $W > 0$. By multiplying (Z, W) with $1/W$, one can assume that $W = 1$. Since E and g are L^0 -linear, the $\operatorname{ess\,sup}$ in (6.4) must be zero, and it follows that $g(Y) = \langle Y, -Z \rangle$ for all $Y \in E$. Moreover, $f(X) \geq \langle X, -Z \rangle$ for all $X \in (L^0)^d$. So $h(X) := \langle X, -Z \rangle$ is the desired extension of g to $(L^0)^d$. □

Theorem 6.7. (Proper separation)

Let C and D be two non-empty σ -stable L^0 -convex subsets of $(L^0)^d$. Then there exists a $Z \in (L^0)^d$ such that

$$\operatorname{ess\,inf}_{X \in C} \langle X, Z \rangle \geq \operatorname{ess\,sup}_{Y \in D} \langle Y, Z \rangle \quad \text{and} \quad \operatorname{ess\,sup}_{X \in C} \langle X, Z \rangle > \operatorname{ess\,inf}_{Y \in D} \langle Y, Z \rangle \quad (6.5)$$

if and only if $0 \notin \operatorname{ri}_A(C - D)$ for all $A \in \mathcal{F}_+$.

Proof. Denote $E = \operatorname{aff}(C - D)$. By Corollary 2.9, $1_A E$ is for all $A \in \mathcal{F}_+$ σ -stable, and therefore, by Proposition 3.5, sequentially closed.

If there exists an $A \in \mathcal{F}_+$ such that $0 \in \operatorname{ri}_A(C - D)$, $1_A E$ is L^0 -linear and there exists an $\varepsilon \in L^0_{++}$ such that $B_\varepsilon^A(0) \cap 1_A E \subseteq 1_A(C - D)$. Suppose there exists $Z \in (L^0)^d$ satisfying (6.5). Then

$$\langle X, Z \rangle \geq 0 \text{ for all } X \in \operatorname{cl}_A(C - D) \quad (6.6)$$

and

$$\langle X, Z \rangle > 0 \text{ on } A \text{ for some } X \in 1_A(C - D). \quad (6.7)$$

One obtains from Corollary 2.12 that $Z = Z_1 + Z_2$ for some $Z_1 \in 1_A E$ and $Z_2 \in (1_A E)^\perp$. It follows from (6.6) that $Z_1 = 0$. But this contradicts (6.7). So (6.5) implies that $0 \notin \operatorname{ri}_A(C - D)$ for all $A \in \mathcal{F}_+$.

Now assume $0 \notin \operatorname{ri}_A(C - D)$ for all $A \in \mathcal{F}_+$. Since E is σ -stable, there exists a largest $B \in \mathcal{F}$ such that $0 \in 1_B E$. If $\mu[B] = 0$, one has $0 \notin 1_A E$ for all $A \in \mathcal{F}_+$, and it follows from Corollary 6.2 that there exists a $Z \in (L^0)^d$ such that $\operatorname{ess\,inf}_{X \in E} \langle X, Z \rangle > 0$, which implies (6.5). If $\mu[B] > 0$, denote $A := \Omega \setminus B$. The same argument as before yields a $Z_0 \in 1_A(L^0)^d$ satisfying (6.6)–(6.7). On the other hand, $1_B E$ is L^0 -linear. So it follows from Corollary 2.11 that there exist disjoint sets $B_1, \dots, B_d \in \mathcal{F}$ satisfying $\bigcup_{i=1}^d B_i = B$ and an orthonormal basis X_1, \dots, X_d of $(L^0)^d$ on B such that $1_{B_i} E = \operatorname{lin}_{B_i} \{X_1, \dots, X_i\}$ for all $i = 1, \dots, d$. For every $i \in \mathcal{I} := \{j = 1, \dots, d : \mu[B_j] > 0\}$ one can apply Theorem 6.4 in the L^0 -linear subset $1_{B_i} E$ to obtain a $Z_i \in 1_{B_i} E$ such that

$$\|Z_i\| > 0 \text{ on } B_i \quad \text{and} \quad \operatorname{ess\,inf}_{X \in C} \langle X, Z_i \rangle \geq \operatorname{ess\,sup}_{Y \in D} \langle Y, Z_i \rangle.$$

Since $0 \notin \operatorname{ri}_A(C - D)$ for all $A \in \mathcal{F}_+$, one has

$$\operatorname{ess\,sup}_{X \in C} \langle X, Z_i \rangle > \operatorname{ess\,inf}_{Y \in D} \langle Y, Z_i \rangle \quad \text{on } B_i.$$

If one sets $Z = 1_A Z_0 + \bigcup_{i \in \mathcal{I}} 1_{B_i} Z_i$, one obtains (6.5), and the proof is complete. \square

7 Properties of L^0 -convex functions

Definition 7.1. Consider a function $f : (L^0)^d \rightarrow L$ and an $X_0 \in (L^0)^d$.

- We call $Y \in (L^0)^d$ an L^0 -subgradient of f at X_0 if

$$f(X_0) \in L^0 \quad \text{and} \quad f(X_0 + X) - f(X_0) \geq \langle X, Y \rangle \quad \text{for all } X \in (L^0)^d.$$

By $\partial f(X_0)$ we denote the set of all L^0 -subgradients of f at X_0 .

- If $f(X_0) \in L^0$ and for some $X \in (L^0)^d$ the limit

$$f'(X_0; X) := \lim_{n \rightarrow \infty} n [f(X_0 + X/n) - f(X_0)]$$

exists a.e. ($+\infty$ and $-\infty$ are allowed as limits), we call it L^0 -directional derivative of f at X_0 in the direction X .

- We say f is L^0 -differentiable at X_0 if $f(X_0) \in L^0$ and there exists a $Y \in (L^0)^d$ such that

$$\frac{f(X_0 + X_n) - f(X_0) - \langle X_n, Y \rangle}{\|X_n\|} \rightarrow 0 \text{ a.e.}$$

for every sequence $(X_n)_{n \in \mathbb{N}}$ in $(L^0)^d$ satisfying $X_n \rightarrow 0$ a.e. and $\|X_n\| > 0$ for all $n \in \mathbb{N}$. If such a Y exists, we call it the L^0 -derivative of f at X_0 and denote it by $\nabla f(X_0)$.

- The L^0 -convex conjugate $f^* : (L^0)^d \rightarrow L$ is given by

$$f^*(Y) := \operatorname{ess\,sup}_{X \in (L^0)^d} \{\langle X, Y \rangle - f(X)\}.$$

- If f is L^0 -convex, we set

$$\operatorname{dom} f := \{X \in (L^0)^d : f(X) < +\infty\}.$$

- By $\operatorname{conv} f$ we denote the largest L^0 -convex function below f and by $\underline{\operatorname{conv}} f$ the largest sequentially lsc L^0 -convex function below f .
- If f is L^0 -convex and satisfies $f(\lambda X) = \lambda f(X)$ for all $\lambda \in L^0_{++}$ and $X \in (L^0)^d$, we call f L^0 -sublinear.
- For every pair $(Y, Z) \in (L^0)^d \times L^0$ we denote by $f^{Y, Z}$ the function from $(L^0)^d$ to L^0 given by $f^{Y, Z}(X) := \langle X, Y \rangle + Z$.

Theorem 7.2. *Let $f : (L^0)^d \rightarrow L$ be an L^0 -convex function and $X_0 \in \operatorname{int}(\operatorname{dom} f)$ such that $f(X_0) \in L^0$. Then $f(X) \in \bar{L}$ for all $X \in (L^0)^d$ and f is sequentially continuous on $\operatorname{int}(\operatorname{dom} f)$.*

Proof. Since $X_0 \in \operatorname{int}(\operatorname{dom} f)$, there exists an $\varepsilon \in L^0_{++}$ such that $V := \max_i f(X_0 \pm \varepsilon e_i) < +\infty$. By L^0 -convexity, one has $f(X) \leq V$ for all $X \in X_0 + U$, where

$$U := \left\{ X \in (L^0)^d : \sum_{i=1}^d |X^i| \leq \varepsilon \right\}.$$

Assume that there exist $X \in (L^0)^d$ and $A \in \mathcal{F}_+$ such that $f(X) = -\infty$ on A . Then one can choose a $Z \in X_0 + U$ and a $\lambda \in L^0$ such that $0 < \lambda \leq 1$ and $X_0 = \lambda X + (1 - \lambda)Z$. It follows that $f(X_0) \leq \lambda f(X) + (1 - \lambda)f(Z) = -\infty$ on A . But this contradicts the assumptions. So $f(X) \in \bar{L}$ for all $X \in (L^0)^d$.

Now pick an $X \in U$ and a $\lambda \in L^0$ such that $0 < \lambda \leq 1$. Then

$$f(X_0 + \lambda X) = f(\lambda(X_0 + X) + (1 - \lambda)X_0) \leq \lambda f(X_0 + X) + (1 - \lambda)f(X_0),$$

and therefore,

$$f(X_0 + \lambda X) - f(X_0) \leq \lambda[f(X_0 + X) - f(X_0)] \leq \lambda(V - f(X_0)).$$

On the other hand,

$$X_0 = \frac{1}{1 + \lambda}(X_0 + \lambda X) + \frac{\lambda}{1 + \lambda}(X_0 - X).$$

So

$$f(X_0) \leq \frac{1}{1 + \lambda}f(X_0 + \lambda X) + \frac{\lambda}{1 + \lambda}f(X_0 - X),$$

which gives

$$f(X_0) - f(X_0 + \lambda X) \leq \lambda[f(X_0 - X) - f(X_0)] \leq \lambda(V - f(X_0)).$$

Hence, we have shown that

$$|f(X) - f(X_0)| \leq \lambda(V - f(X_0)) \quad \text{for all } X \in X_0 + \lambda U.$$

Let $(X_n)_{n \in \mathbb{N}}$ be a sequence in $(L^0)^d$ converging a.e. to X_0 . For every $k \in \mathbb{N}$, the sets

$$A_m^k := \bigcap_{n \geq m} \{X_n - X_0 \in U/k\}$$

are increasing in m with $\bigcup_{m \geq 1} A_m^k = \Omega$. By Lemma 4.3, f is stable. Therefore,

$$|f(X_n) - f(X_0)| \leq (V - f(X_0))/k \quad \text{for all } n \geq m \quad \text{on } A_m^k,$$

and one obtains

$$\mu \left[\bigcup_{k \geq 1} \bigcap_{m \geq 1} \bigcup_{n \geq m} \{|f(X_n) - f(X_0)| > (V - f(X_0))/k\} \right] = 0.$$

So $f(X_n) \rightarrow f(X_0)$ a.e., and the theorem follows. \square

As an immediate consequence of Theorem 7.2 one obtains the following

Corollary 7.3. *An L^0 -convex function $f : (L^0)^d \rightarrow \bar{L}$ is sequentially continuous on $\text{int}(\text{dom } f)$.*

Theorem 7.4. *Let $f : (L^0)^d \rightarrow \bar{L}$ be an L^0 -convex function and $X_0 \in \text{ri}(\text{dom } f)$. Then $\partial f(X_0) \neq \emptyset$. In particular, if $f(X) \in L^0$ for all $X \in (L^0)^d$, then $\partial f(X_0) \neq \emptyset$ for all $X \in (L^0)^d$.*

Proof. By Lemma 4.3, f is stable. Therefore,

$$C := \{(X, V) \in (L^0)^d \times L^0 : f(X) \leq V\}$$

is an L^0 -convex, σ -stable subset of $(L^0)^d \times L^0$. Since $(X_0, f(X_0) + 1)$ is in C , one has $(0, 0) \notin \text{ri}_A(C - (X_0, f(X_0)))$ for all $A \in \mathcal{F}_+$. So it follows from Theorem 6.7 that there exists $(Y, Z) \in (L^0)^d \times L^0$ such that

$$\text{ess inf}_{(X, V) \in C} \{\langle X, Y \rangle + VZ\} \geq \langle X_0, Y \rangle + f(X_0)Z \quad (7.1)$$

and

$$\text{ess sup}_{(X, V) \in C} \{\langle X, Y \rangle + VZ\} > \langle X_0, Y \rangle + f(X_0)Z. \quad (7.2)$$

(7.1) implies that $Z \geq 0$. Now assume there exists an $A \in \mathcal{F}_+$ such that $1_A Z = 0$. Then since $X_0 \in \text{ri}(\text{dom } f)$, (7.2) contradicts (7.1). So one must have $Z > 0$, and by multiplying (Y, Z) with $1/Z$, one can assume $Z = 1$. It follows from (7.1) that

$$\text{ess inf}_{X \in \text{dom } f} \{\langle X, Y \rangle + f(X)\} = \langle X_0, Y \rangle + f(X_0),$$

which shows that $-Y$ is an L^0 -subgradient of f at X_0 . □

Lemma 7.5. *Let $f, g : (L^0)^d \rightarrow L$ be functions such that $f \geq g$. Then the following hold:*

- (i) f^* is sequentially lsc and L^0 -convex;
- (ii) $f^*(Y) \geq \langle X, Y \rangle - f(X)$ for all $X, Y \in (L^0)^d$;
- (iii) $Y \in \partial f(X)$ if and only if $f(X) \in L^0$ and $f^*(Y) = \langle X, Y \rangle - f(X)$;
- (iv) $f^* \leq g^*$ and $f^{**} \geq g^{**}$;
- (v) $f \geq f^{**}$ and $f^* = f^{***}$.

Proof. To prove (i) let $(Y_n)_{n \in \mathbb{N}}$ be a sequence in $(L^0)^d$ converging a.e. to some $Y \in (L^0)^d$. Then

$$\begin{aligned} \liminf_{n \rightarrow \infty} f^*(Y_n) &= \sup_{m \geq 1} \inf_{n \geq m} \text{ess sup}_{X \in (L^0)^d} \{\langle X, Y_n \rangle - f(X)\} \\ &\geq \text{ess sup}_{X \in (L^0)^d} \sup_{m \geq 1} \inf_{n \geq m} \{\langle X, Y_n \rangle - f(X)\} \\ &= \text{ess sup}_{X \in (L^0)^d} \{\langle X, Y \rangle - f(X)\} = f^*(Y). \end{aligned}$$

Hence, f^* is sequentially lsc. To show that it is L^0 -convex, choose $Y, Z \in (L^0)^d$ and $\lambda \in L^0$ such that $0 \leq \lambda \leq 1$. Then, $\lambda f^*(Y) + (1 - \lambda)f^*(Z) \geq \langle X, \lambda Y + (1 - \lambda)Z \rangle - f(X)$ for all $X \in (L^0)^d$ and therefore, $\lambda f^*(Y) + (1 - \lambda)f^*(Z) \geq f^*(\lambda Y + (1 - \lambda)Z)$. (ii) is immediate from the definition of f^* . Now assume that $f(X) \in L^0$. For any $X' \in (L^0)^d$, $f(X') - f(X) \geq \langle X' - X, Y \rangle$ is equivalent to $\langle X, Y \rangle - f(X) \geq \langle X', Y \rangle - f(X')$. This shows (iii). (iv) is clear. From (ii) one obtains that $f(X) \geq \langle X, Y \rangle - f^*(Y)$ for all $X, Y \in (L^0)^d$. So $f \geq f^{**}$. The same inequality applied to f^* gives $f^* \geq f^{***}$. On the other hand, we know from (iv) that $f^* \leq f^{***}$. This proves (v). □

Lemma 7.6. *Let $f : (L^0)^d \rightarrow \bar{L}$ be a sequentially lsc L^0 -convex function. Then one has for all $X \in (L^0)^d$,*

$$f(X) = \text{ess sup} \{f^{Y, Z}(X) : (Y, Z) \in (L^0)^d \times L^0, f \geq f^{Y, Z}\}.$$

Proof. Note that the set

$$\mathcal{A} := \{A \in \mathcal{F} : \text{there exists an } X \in (L^0)^d \text{ such that } 1_A f(X) \in L^0\}$$

is directed upwards. Therefore, there exists an increasing sequence A_n in \mathcal{A} with corresponding X_n , $n \in \mathbb{N}$, such that $A_n \uparrow A := \text{ess sup } \mathcal{A}$ a.e. Set

$$X_0 := 1_{A_1 \cup A^c} X_1 + \sum_{n \geq 2} 1_{A_n \setminus A_{n-1}} X_n.$$

By Lemma 4.3, f is stable. Hence, $f(X_0) < +\infty$ on A , and $f(X) = +\infty$ on A^c for all $X \in (L^0)^d$. The lemma can be proved on A and A^c separately, and on A^c it is obvious. Therefore, we can assume $A = \Omega$. Then $\text{dom } f \neq \emptyset$, and it follows that

$$C := \{(X, V) \in \text{dom } f \times L^0 : f(X) \leq V\}$$

is a non-empty sequentially closed L^0 -convex subset of $(L^0)^d \times L^0$. Choose a pair $(U, W) \in (L^0)^d \times L^0$ such that $1_A(U, W) \notin 1_{A^c} C$ for all $A \in \mathcal{F}_+$. By Corollary 6.2, there exists $(Y, Z) \in (L^0)^d \times L^0$ such that

$$I := \inf_{(X, V) \in C} \{\langle X, Y \rangle + VZ\} > \langle U, Y \rangle + WZ.$$

It follows that $Z \geq 0$. On the set $B := \{Z > 0\}$ one can multiply (Y, Z) with $1/Z$ and assume $Z = 1$. Then one obtains that on B ,

$$f(X) \geq f^{-Y, I}(X) \quad \text{for all } X \in (L^0)^d \quad \text{and} \quad f^{-Y, I}(U) > W.$$

On B^c one has $\lambda := I - \langle U, Y \rangle > 0$. Pick a $U' \in \text{dom } f$. Since $1_A(U', f(U') - 1) \notin 1_{A^c} C$ for all $A \in \mathcal{F}_+$, one obtains from Corollary 6.2 that there exists a pair $(Y', Z') \in (L^0)^d \times L^0$ such that

$$I' := \inf_{(X, V) \in C} \{\langle X, Y' \rangle + VZ'\} > \langle U', Y' \rangle + (f(U') - 1)Z'.$$

Since $U' \in \text{dom } f$, one must have $Z' > 0$. By multiplying with $1/Z'$, one can assume $Z' = 1$. Now choose a $\delta \in 1_{B^c} L^0_+$ such that

$$\delta > \frac{1}{\lambda}(W + \langle U, Y' \rangle - I)^+ \quad \text{on } B^c$$

and set $Y'' := \delta Y + Y'$. Then, on B^c ,

$$I'' := \inf_{(X, V) \in C} (\langle X, Y'' \rangle + V) \geq \delta I + I' = \delta \lambda + \delta \langle U, Y \rangle + I' > \langle U, Y'' \rangle + W.$$

So on B^c , one has

$$f(X) \geq f^{-Y'', I''}(X) \quad \text{for all } X \in (L^0)^d \quad \text{and} \quad f^{-Y'', I''}(U) > W.$$

Now define $(\hat{Y}, \hat{I}) := 1_B(-Y, I) + 1_{B^c}(-Y'', I'')$. Then

$$f(X) \geq f^{\hat{Y}, \hat{I}}(X) \quad \text{for all } X \in (L^0)^d \quad \text{and} \quad f^{\hat{Y}, \hat{I}}(U) > W.$$

This proves the lemma. □

Theorem 7.7. (Conditional version of the Fenchel–Moreau theorem)

Let $f : (L^0)^d \rightarrow \bar{L}$ be a function such that $\underline{\text{conv}} f$ takes values in \bar{L} . Then $\underline{\text{conv}} f = f^{**}$. In particular, if f is sequentially lsc and L^0 -convex, then $f = f^{**}$.

Proof. We know from Lemma 7.5 that f^{**} is a sequentially lsc L^0 -convex minorant of f . So $\underline{\text{conv}} f \geq f^{**}$. On the other hand, it follows from Lemma 7.6 that

$$\underline{\text{conv}} f = \text{ess sup} \{f^{Y, Z}(X) : (Y, Z) \in (L^0)^d \times L^0, \underline{\text{conv}} f \geq f^{Y, Z}\},$$

and it can easily be checked that $(f^{Y, Z})^{**} = f^{Y, Z}$ for all $(Y, Z) \in (L^0)^d \times L^0$. So one obtains from Lemma 7.5 that $f^{**} \geq (f^{Y, Z})^{**} = f^{Y, Z}$ for every pair $(Y, Z) \in (L^0)^d \times L^0$ satisfying $f \geq f^{Y, Z}$. This shows that $f^{**} \geq \underline{\text{conv}} f$. □

Lemma 7.8. *Let $f : (L^0)^d \rightarrow L$ be an L^0 -convex function and $X_0 \in (L^0)^d$ such that $f(X_0) \in L^0$. Then $f'(X_0; X)$ exists for all $X \in (L^0)^d$, $f'(X_0, 0) = 0$ and $f'(X_0; \cdot)$ is L^0 -sublinear. Moreover, $\partial f(X_0) = \partial g(0)$, where $g(X) := f'(X_0; X)$.*

Proof. It follows from L^0 -convexity that for every $X \in (L^0)^d$, $n[f(X_0 + X/n) - f(X_0)]$ is decreasing in n . This implies that $f'(X_0; X)$ exists. $f'(X_0; 0) = 0$ is clear. That $f'(X_0; \cdot)$ is L^0 -sublinear and $\partial f(X_0) = \partial g(0)$ are straightforward to check. \square

Lemma 7.9. *Let $f : (L^0)^d \rightarrow \bar{L}$ be a sequentially lsc L^0 -sublinear function. If there exists an $X_0 \in (L^0)^d$ such that $f(X_0) \in L^0$, then $\partial f(0) \neq \emptyset$ and $f(X) = \text{ess sup}_{Y \in \partial f(0)} \langle X, Y \rangle$ for all $X \in (L^0)^d$. In particular, $f(0) = 0$.*

Proof. By Theorem 7.7, one has $f = f^{**}$. This implies that the set

$$C := \{Y \in (L^0)^d : \langle X, Y \rangle \leq f(X) \text{ for all } X \in (L^0)^d\}$$

is non-empty and $f(X) = \text{ess sup}_{Y \in C} \langle X, Y \rangle$. It follows that $f(0) = 0$ and $\partial f(0) = C$. This proves the lemma. \square

Theorem 7.10. *Let $f : (L^0)^d \rightarrow \bar{L}$ be an L^0 -convex function. Assume there exist $X_0 \in (L^0)^d$ and $V \in L_+^0$ such that $f(X_0) \in L^0$ and*

$$f(X_0 + X) \geq f(X_0) - V\|X\| \quad \text{for all } X \in (L^0)^d. \quad (7.3)$$

Then there exists a $Y \in \partial f(X_0)$ such that $\|Y\| \leq V$.

Proof. Denote $g(X) := f'(X_0; X)$. Then $h = \text{conv} g$ is a sequentially lsc L^0 -sublinear function which by (7.3), satisfies

$$h(X) \geq -V\|X\| \quad \text{for all } X \in (L^0)^d. \quad (7.4)$$

It follows that $h(0) = 0$ and $\partial h(0) \subseteq \partial g(0) = \partial f(X_0)$. Since $\partial h(0)$ and

$$B^V(0) := \{Y \in (L^0)^d : \|Y\| \leq V\}$$

are L^0 -convex and sequentially closed, they are both σ -stable. Therefore, there exists a largest set $A \in \mathcal{F}$ such that $1_A \partial h(0) \cap 1_A B^V(0)$ is non-empty. Assume that $A^c \in \mathcal{F}_+$. Then, if one restricts attention to A^c and assumes $\Omega = A^c$, the sets $\partial h(0)$ and $B^V(0)$ satisfy the assumptions of Corollary 6.2. So there exists a $Z \in (L^0)^d$ such that

$$-V\|Z\| = \text{ess inf}_{Y \in B^V(0)} \langle Y, Z \rangle > \text{ess sup}_{Y \in \partial h(0)} \langle Y, Z \rangle.$$

But by Lemma 7.9, one has $h(Z) = \text{ess sup}_{Y \in \partial h(0)} \langle Y, Z \rangle$, and one obtains a contradiction to (7.4). It follows that $A = \Omega$, which proves the theorem. \square

Theorem 7.11. *Let $f : (L^0)^d \rightarrow \bar{L}$ be an L^0 -convex function and X_0 in $(L^0)^d$ such that $f(X_0) \in L^0$. Assume that $\partial f(X_0) = \{Y\}$ for some $Y \in (L^0)^d$. Then f is L^0 -differentiable at X_0 with $\nabla f(X_0) = Y$.*

Proof. By Lemma 7.8, one has $\partial g(0) = \{Y\}$ for the L^0 -sublinear function $g(X) := f'(X_0; X)$. It follows that

$$g^*(Z) = 1_{\{Z \neq Y\}}(+\infty) \quad \text{and} \quad g^{**}(X) = \langle X, Y \rangle. \quad (7.5)$$

Set

$$\mathcal{A} := \{A \in \mathcal{F} : \text{there exists an } X \in (L^0)^d \text{ such that } g(X) = +\infty \text{ on } A\}.$$

By Lemma 4.3, g is stable. Therefore, there exists a sequence $(A_n)_{n \in \mathbb{N}}$ in \mathcal{A} with corresponding X_n such that $A_n \uparrow A := \text{ess sup } \mathcal{A}$. The element

$$X_0 := 1_{A_1 \cup A^c} X_1 + \sum_{n \geq 2} 1_{A_n \setminus A_{n-1}} X_n$$

satisfies $g(X_0) = +\infty$ on A . We want to show that $\mu[A] = 0$. So let us assume $\mu[A] > 0$. If one replaces Ω with A , one has $0 \notin 1_B(\text{dom } g - X_0)$ for all $B \in \mathcal{F}_+$. By Theorem 6.4, there exists a $Z \in (L^0)^d$ such that

$$\|Z\| > 0 \quad \text{and} \quad \text{ess inf}_{X \in \text{dom } g} \langle X, Z \rangle \geq \langle X_0, Z \rangle.$$

Define the sequentially lsc L^0 -convex function $h : (L^0)^d \rightarrow \bar{L}$ as follows:

$$h(X) := \langle X, Y \rangle 1_{\{\langle X, Z \rangle \geq \langle X_0, Z \rangle\}} + \infty 1_{\{\langle X, Z \rangle < \langle X_0, Z \rangle\}}.$$

Then $g \geq h$ and $h(X) = +\infty$ for all $X \in (L^0)^d$ satisfying $\langle X, Z \rangle < \langle X_0, Z \rangle$. It follows that $\text{conv}g(X) = +\infty$ for all $X \in (L^0)^d$ satisfying $\langle X, Z \rangle < \langle X_0, Z \rangle$. Moreover, since $Y \in \partial g(0)$, g fulfills the assumptions of Theorem 7.7, and one obtains $\text{conv}g = g^{**}$, contradicting (7.5). So one must have $\mu[A] = 0$, or in other words, $g(X) \in L^0$ for all $X \in (L^0)^d$. It follows from Theorem 7.2 that g is sequentially continuous, and therefore, $g(X) = g^{**}(X) = \langle X, Y \rangle$ for all $X \in (L^0)^d$.

Now let $(X_n)_{n \in \mathbb{N}}$ be a sequence in $(L^0)^d$ such that $X_n \rightarrow 0$ a.e. and $\|X_n\| > 0$ for all n . Denote $\|X_n\|_1 := \sum_{i=1}^d |X_n^i|$ and notice that there exists a constant $c > 0$ such that $\|X_n\|_1 \leq c\|X_n\|$ for all n . Since $g(X) = \langle X, Y \rangle$, one has for all $i = 1, \dots, d$,

$$\frac{f(X_0 \pm \|X_n\|_1 e_i) - f(X_0)}{\|X_n\|_1} \rightarrow \pm Y^i \quad \text{a.e.}$$

Therefore,

$$\begin{aligned} & \frac{f(X_0 + X_n) - f(X_0) - \langle X_n, Y \rangle}{\|X_n\|} \leq c \frac{f(X_0 + X_n) - f(X_0) - \langle X_n, Y \rangle}{\|X_n\|_1} \\ & \leq c \sum_{i=1}^d \frac{|X_n^i|}{\|X_n\|_1} \left\{ \frac{f(X_0 + \|X_n\|_1 \text{sign}(X_n^i) e_i) - f(X_0)}{\|X_n\|_1} - \text{sign}(X_n^i) Y^i \right\} \rightarrow 0 \quad \text{a.e.} \end{aligned}$$

□

8 Inf-convolution

Definition 8.1. We define the inf-convolution of finitely many functions $f_j : (L^0)^d \rightarrow \bar{L}$, $j = 1, \dots, n$, by

$$\square_{j=1}^n f_j(X) := \text{ess inf}_{X_1 + \dots + X_n = X} \sum_{j=1}^n f_j(X_j).$$

Lemma 8.2. If f_j , $j = 1, \dots, n$, are L^0 -convex functions from $(L^0)^d$ to \bar{L} , then $\square_{j=1}^n f_j$ is L^0 -convex too.

Proof. Denote $f = \square_{j=1}^n f_j$. Choose $X, Y \in (L^0)^d$ and $V, W \in \bar{L}$ such that $f(X) \leq V$ and $f(Y) \leq W$. Let $\varepsilon \in L_{++}^0$ and $\lambda \in L^0$ such that $0 \leq \lambda \leq 1$. By Lemma 4.3, the functions f_j are stable. Therefore, the family $\left\{ \sum_j f_j(X_j) : \sum_j X_j = X \right\}$ is directed downwards. So there exist sequences X_j^k , $k \in \mathbb{N}$, such that $\sum_j X_j^k = X$ and $\sum_j f_j(X_j^k)$ decreases to $f(X)$ a.e. It follows that the sets $A_k := \left\{ \sum_j f_j(X_j^k) \leq V + \varepsilon \right\}$ increase to Ω as $k \rightarrow \infty$. So for every $j = 1, \dots, n$,

$$X_j := \sum_{k \geq 1} 1_{A_k \setminus A_{k-1}} X_j^k, \quad \text{where } A_0 := \emptyset.$$

defines an element in $(L^0)^d$ such that $\sum_{j=1}^n X_j = X$ and $\sum_{j=1}^n f(X_j) \leq V + \varepsilon$. Analogously, there exist $Y_j \in (L^0)^d$, $j = 1, \dots, n$, such that $\sum_{j=1}^n Y_j = Y$ and $\sum_{j=1}^n f(Y_j) \leq W + \varepsilon$. Set $Z_j = \lambda X_j + (1 - \lambda) Y_j$. Then $Z := \sum_{j=1}^n Z_j = \lambda X + (1 - \lambda) Y$ and

$$f(Z) \leq \sum_{j=1}^n f_j(Z_j) \leq \sum_{j=1}^n \lambda f_j(X_j) + (1 - \lambda) f_j(Y_j) \leq \lambda V + (1 - \lambda) W + \varepsilon.$$

It follows that $f(Z) \leq \lambda f(X) + (1 - \lambda) f(Y)$. □

Lemma 8.3. Let $f_j : (L^0)^d \rightarrow \bar{L}$, $j = 1, \dots, n$, be L^0 -convex functions and denote $f = \square_{j=1}^n f_j$. Assume $f(X_0) = \sum_{j=1}^n f_j(X_j) < +\infty$ for some $X_j \in (L^0)^d$ summing up to X_0 . If $X_1 \in \text{int}(\text{dom } f_1)$, then $f(X) \in \bar{L}$ for all $X \in (L^0)^d$, $X_0 \in \text{int}(\text{dom } f)$ and f is sequentially continuous on $\text{int}(\text{dom } f)$.

Proof. By definition of f , one has

$$f(X_0 + X) - f(X_0) \leq f_1(X_1 + X) + \sum_{j=2}^n f_j(X_j) - \sum_{j=1}^n f_j(X_j) = f_1(X_1 + X) - f_1(X_1)$$

for all $X \in (L^0)^d$. This shows that $X_0 \in \text{int}(\text{dom } f)$. Since $f(X_0) = \sum_{j=1}^n f_j(X_j) \in L^0$, the rest of the lemma follows from Theorem 7.2. \square

Lemma 8.4. Consider functions $f_j : (L^0)^d \rightarrow \bar{L}$, $j = 1, \dots, n$, and denote $f = \square_{j=1}^n f_j$. Assume $f(X_0) = \sum_{j=1}^n f_j(X_j) < +\infty$ for some $X_j \in (L^0)^d$ summing up to X_0 . Then $\partial f(X_0) = \bigcap_{j=1}^n \partial f_j(X_j)$.

Proof. Assume $Y \in \partial f(X_0)$ and $X \in (L^0)^d$. Then

$$f_1(X_1 + X) - f_1(X_1) = f_1(X_1 + X) + \sum_{j=2}^n f_j(X_j) - \sum_{j=1}^n f_j(X_j) \geq f(X_0 + X) - f(X_0) \geq \langle X, Y \rangle.$$

Hence $Y \in \partial f_1(X_1)$, and by symmetry, $\partial f(X_0) \subseteq \bigcap_{j=1}^n \partial f_j(X_j)$. On the other hand, if $Y \in \bigcap_{j=1}^n \partial f_j(X_j)$ and $X \in (L^0)^d$, choose Z_j such that $\sum_{j=1}^n Z_j = X_0 + X$. Then

$$\sum_{j=1}^n f_j(Z_j) \geq \sum_{j=1}^n f_j(X_j) + \langle Z_j - X_j, Y \rangle = \sum_{j=1}^n f_j(X_j) + \langle X, Y \rangle.$$

So $f(X_0 + X) - f(X_0) \geq \langle X, Y \rangle$, and the lemma follows. \square

Lemma 8.5. Let $f_j : (L^0)^d \rightarrow \bar{L}$, $j = 1, \dots, n$, be L^0 -convex functions and denote $f = \square_{j=1}^n f_j$. Assume $f(X_0) = \sum_j f_j(X_j) < +\infty$ for some $X_j \in (L^0)^d$ summing up to X_0 and f_1 is L^0 -differentiable at X_1 . Then f is L^0 -differentiable at X_0 with $\nabla f(X_0) = \nabla f_1(X_1)$.

Proof. One has

$$f(X_0 + X) - f(X_0) \leq f_1(X_1 + X) + \sum_{j=2}^n f_j(X_j) - \sum_{j=1}^n f_j(X_j) = f_1(X_1 + X) - f_1(X_1)$$

for all $X \in (L^0)^d$. It follows that the L^0 -directional derivative $g(X) := f'(X_0; X)$ satisfies

$$g(X) \leq f'_1(X_1; X) = \langle X, \nabla f_1(X_1) \rangle$$

for all $X \in (L^0)^d$. But by Lemma 8.2, f is L^0 -convex. It follows that g is L^0 -sublinear, and therefore, $g(X) = \langle X, \nabla f_1(X_1) \rangle$. This implies that $\partial f(X_0) = \partial g(0) = \{\nabla f_1(X_1)\}$. Now the lemma follows from Theorem 7.11. \square

Lemma 8.6. Consider functions $f_j : (L^0)^d \rightarrow \bar{L}$, $j = 1, \dots, n$. Then $(\square_{j=1}^n f_j)^* = \sum_{j=1}^n f_j^*$, where the sum is understood to be $-\infty$ if at least one of the terms is $-\infty$.

Proof.

$$(\square_{j=1}^n f_j)^*(Y) = \text{ess sup}_X \{ \langle X, Y \rangle - \square_{j=1}^n f_j(X) \} = \text{ess sup}_{X_1, \dots, X_n} \sum_{j=1}^n \{ \langle X_j, Y \rangle - f_j(X_j) \} = \sum_{j=1}^n f_j^*(Y).$$

\square

References

- [1] Cheridito, P., Hu, Y. (2011). Optimal consumption and investment in incomplete markets with general constraints. *Stochastics and Dynamics* 11(2), 283–299.
- [2] Cheridito, P., Horst, U., Kupper, M., Pirvu, T. (2014). Equilibrium pricing in incomplete markets under translation invariant preferences. SSRN Preprint.

- [3] Cheridito, P., Stadjje, M. (2012). BS Δ Es and BSDEs with non-Lipschitz drivers: comparison, convergence and robustness. Forthcoming in Bernoulli.
- [4] Filipović, D., Kupper, M., Vogelpoth, N. (2009). Separation and Duality in Locally L^0 -Convex Modules. *Journal of Functional Analysis* 256, 3996–4029.
- [5] Föllmer, H., Schied, A. (2004). *Stochastic Finance, An Introduction in Discrete Time*. 2nd Edition. de Gruyter Studies in Mathematics 27.
- [6] Guo, T. (2010). Relations between some basic results derived from two kinds of topologies for a random locally convex module. *Journal of Functional Analysis* 258, 3024–3047.
- [7] Guo, T. (2011). Recent progress in random metric theory and its applications to conditional risk measures. *Science China Mathematics* 54(4), 633–660.
- [8] Kabanov, Y., Stricker, Ch. (2001) A teacher’s note on no-arbitrage criteria. *Séminaire de probabilités*, 35, 149–152.
- [9] Kupper, M., Vogelpoth, N. (2009). Complete L^0 -normed modules and automatic continuity of monotone convex functions. Preprint.
- [10] Neveu, J. (1975). *Discrete-Parameter Martingales*. North-Holland Publishing Company Amsterdam, Oxford.