

**REAL ALGEBRAIC GEOMETRY LECTURE NOTES**  
**(07: 10/11/09 - BEARBEITET 19/11/2018)**

SALMA KUHLMANN

CONTENTS

1. Sturm's Theorem	1
--------------------	---

Let  $R$  be a real closed field.

1. STURM'S THEOREM

**Definition 1.1.**

- (i) Let  $f \in R[x]$  be a non-constant polynomial,  $\deg(f) \geq 1$ . The **Sturm sequence** of  $f$  is defined recursively as the sequence  $(f_0, \dots, f_r)$  of polynomials in  $R[x]$  such that:

$$\begin{aligned} f_0 &:= f, & f_1 &:= f' & \text{and} \\ f_0 &= f_1 q_1 - f_2 \\ f_1 &= f_2 q_2 - f_3 \\ &\dots \\ f_{i-1} &= f_i q_i - f_{i+1} \\ &\dots \\ f_{r-2} &= f_{r-1} q_{r-1} - f_r \\ f_{r-1} &= f_r q_r, \end{aligned}$$

where  $f_i, q_i \in R[x]$ ,  $f_i \neq 0$  and  $\deg(f_i) < \deg(f_{i-1})$   $r, f_i, q_i$  uniquely determined.

- (ii) Let  $x \in R$ . Set

$$V_f(x) := \text{Var}(f_0(x), \dots, f_r(x)).$$

We recall that after we have removed all zero's by the sequence  $(c_1, \dots, c_n)$ , we defined  $\text{Var}(c_1, \dots, c_n)$  as the number of changes of sign in  $(c_1, \dots, c_n)$ , i.e.

$$\text{Var}(c_1, \dots, c_n) = |\{i \in \{1, \dots, n\} : c_i c_{i+1} < 0\}|.$$

**Theorem 1.2.** (*Sturm 1829*). Let  $a, b \in R$ ,  $a < b$ ,  $f(a)f(b) \neq 0$ . Then

$$|\{c : a \leq c \leq b, f(c) = 0\}| = V_f(a) - V_f(b).$$

*Proof.* For the proof we study the function  $V_f(x)$ ,  $x \in R$ , locally constant except around finitely many roots for  $f_0, \dots, f_r$ .

- (1) Suppose  $\gcd(f_0, f_1) = 1$ .  
(2) Hilfslemma (ÜA)  $\exists \delta$  such that

$$|x - c| < \delta \Rightarrow \text{sign}(f_0(x)f_1(x)) = \text{sign}(x - c) = \begin{cases} -1 & \text{if } x < c \\ 0 & \text{if } x = c \\ 1 & \text{if } x > c. \end{cases}$$

- (3)  $\forall i \in \{1, \dots, r-1\}$ :  $\gcd(f_{i-1}, f_i) = 1$  and

$$f_{i-1} = q_i f_i - f_{i+1}, \quad \text{with } f_{i+1} \neq 0.$$

So if  $f_i(c) = 0$  then

$$f_{i-1}(c)f_{i+1}(c) < 0.$$

- (4) Let  $f_i(c) = 0$  for  $i \in \{0, \dots, r-1\}$ . Then  $f_{i+1}(c) \neq 0$  (so  $\text{sign}(f_{i+1}(c)) = \pm 1$ ).

We shall now compare for  $f_i(c) = 0$ ,  $i \in \{0, \dots, r-1\}$

$$\text{sign}(f_i(x)) \quad \text{sign}(f_{i+1}(x))$$

for  $|x - c| < \delta$  and count.

We first examine the case  $i = 0$ .

Observe that  $\text{sign}(f_1(x)) \neq 0 \forall x$  such that  $|x - c| < \delta$  because of Hilfslemma. So in particular  $\text{sign}(f_1(x))$  is constant for  $|x - c| < \delta$  and it is equal to  $\text{sign}(f_1(c))$ :

	$x \rightarrow c_-$	$x = c$	$x \rightarrow c_+$
$f_0(x)$	$-\text{sign}(f_1(c))$	0	$\text{sign}(f_1(c))$
$f_1(x)$	$\text{sign}(f_1(c))$	$\text{sign}(f_1(c))$	$\text{sign}(f_1(c))$
contribution to $V_f(x)$	1	0	0

Now consider  $i \in \{1, \dots, r-1\}$  and use (3), i.e.

$$f_i(d) = 0 \implies f_{i-1}(d)f_{i+1}(d) < 0:$$

	$x \rightarrow d_-$	$x = d$	$x \rightarrow d_+$
$f_{i-1}(x)$	$-\text{sign}(f_{i+1}(d))$	$-\text{sign}(f_{i+1}(d))$	$-\text{sign}(f_{i+1}(d))$
$f_i(x)$		0	
$f_{i+1}(x)$	$\text{sign}(f_{i+1}(d))$	$\text{sign}(f_{i+1}(d))$	$\text{sign}(f_{i+1}(d))$
contribution to $V_f(x)$	1	1	1

Therefore for  $a < b$ ,  $V_f(a) - V_f(b)$  is the number of roots of  $f$  in  $]a, b[$ .

Let us consider now the general case. Set

$$g_i := f_i/f_r \quad i = 0, \dots, r.$$

The sequence of polynomials  $(g_0, \dots, g_r)$  satisfies the previous conditions (1) – (4). We can conclude by noticing that:

$$(i) \text{Var}(g_0(x), \dots, g_r(x)) = \text{Var}(f_0(x), \dots, f_r(x)) \text{ (because } f_i(x) = f_r(x)g_i(x)\text{),}$$

(ii)  $f = f_0$  and  $g_0 = f/f_r$  have the same zeros ( $f_r = \gcd(f, f')$ , so  $g = f/f_r$  has only simple roots, whereas  $f$  has roots with multiplicities.)

□

For  $i = 0, \dots, r$  set  $d_i := \deg(f_i)$  and  $\varphi_i :=$  the leading coefficient of  $f_i$ .  
Set

$$V_f(-\infty) := \text{Var}((-1)^{d_0}\varphi_0, (-1)^{d_1}\varphi_1, \dots, (-1)^{d_r}\varphi_r)$$

$$V_f(+\infty) := \text{Var}(\varphi_0, \varphi_1, \dots, \varphi_r).$$

Then we have:

**Corollary 1.3.** *The number of distinct roots of  $f$  is  $V_f(-\infty) - V_f(+\infty)$ .*