

**REAL ALGEBRAIC GEOMETRY LECTURE NOTES**  
**(04: 29/10/09)**

SALMA KUHLMANN

CONTENTS

|                                |   |
|--------------------------------|---|
| 1. Ordering extensions         | 1 |
| 2. Quadratic extensions        | 1 |
| 3. Odd degree field extensions | 2 |
| 4. Real closed fields          | 3 |

1. ORDERING EXTENSIONS

**Definition 1.1.** Let  $L/K$  be a field extension and  $P$  an ordering on  $K$ .

An ordering  $Q$  of  $L$  is said to be an **extension** (*Fortsetzung*) of  $P$  if  $P \subset Q$  (equivalently  $Q \cap K = P$ ).

**Definition 1.2.** Let  $L/K$  be a field extension and  $P$  an ordering on  $K$ . We define

$$T_L(P) := \left\{ \sum_{i=1}^n p_i y_i^2 : n \in \mathbb{N}, p_i \in P, y_i \in L \right\}.$$

**Remark 1.3.** Let  $L/K$  be a field extension and  $P$  an ordering on  $K$ .

Then  $T_L(P)$  is the smallest preordering of  $L$  containing  $P$ .

**Corollary 1.4.** Let  $L/K$  be a field extension and  $P$  an ordering on  $K$ .

Then  $P$  has an extension to an ordering  $Q$  of  $L$  if and only if  $T_L(P)$  is a proper preordering (i.e. if and only if  $-1 \notin T_L(P)$ ).

2. QUADRATIC EXTENSIONS

**Theorem 2.1.** Let  $K$  be a field,  $a \in K$  and define  $L := K(\sqrt{a})$ . Then an ordering  $P$  of  $K$  extends to an ordering  $Q$  of  $L$  if and only if  $a \in P$ .

*Proof.*

( $\Rightarrow$ ) Assume  $Q$  is an extension of  $P$ , then  $a = (\sqrt{a})^2 \in Q \cap K = P$ .

( $\Leftarrow$ ) Let  $a \in P$  (without loss of generality we can assume  $L \neq K$  and  $\sqrt{a} \notin K$ ). We show that  $T_L(P)$  is a proper preordering (and then the thesis follows by Corollary 1.4).

If not, there is  $n \in \mathbb{N}$  and there are  $x_1, \dots, x_n, y_1, \dots, y_n \in K$ ,  $p_1, \dots, p_n \in P$  such that

$$\begin{aligned}
-1 &= \sum_{i=1}^n p_i (x_i + y_i \sqrt{a})^2 \\
&= \sum_{i=1}^n p_i (x_i^2 + ay_i^2 + 2x_i y_i \sqrt{a}).
\end{aligned}$$

On the other hand  $-1 \in K$ , and since every  $x \in K(\sqrt{a})$  can be written in a unique way as  $x = k_1 + k_2 \sqrt{a}$  with  $k_1, k_2 \in K$ , it follows that

$$-1 = \sum_{i=1}^n p_i (x_i^2 + ay_i^2) \in P,$$

contradiction. □

### 3. ODD DEGREE FIELD EXTENSIONS

**Theorem 3.1.** *Let  $L/K$  be a field extension such that  $[L : K]$  is finite and odd. Then every ordering of  $K$  extends to an ordering of  $L$ .*

*Proof.* Otherwise, let  $n \in \mathbb{N}$  the minimal odd degree of a field extension for which the theorem fails.

Let  $L/K$  be a finite field extension such that  $[L : K] = n$  and let  $P$  be an ordering of  $K$  not extending to an ordering of  $L$ .

Since  $\text{char}(K) = 0$  Primitive Element Theorem applies and there is some  $\alpha \in L \setminus K$  such that

$$L = K(\alpha) \cong K[x]/(f),$$

where  $f$  is the minimal polynomial of  $\alpha$  over  $K$ . Therefore  $\deg(f) = n$ ,  $f(\alpha) = 0$  and for every  $g(x) \in K[x]$  such that  $\deg(g) < n$ , we have  $g(\alpha) \neq 0$ .

By Corollary 1.4,  $-1 \in T_L(P)$ , so

$$1 + \sum_{i=1}^s p_i y_i^2 = 0,$$

where  $\forall i = 1, \dots, s$   $p_i \in P$ ,  $p_i \neq 0$ ,  $y_i \in L$ ,  $y_i \neq 0$ . Define

$$y_i = g_i(\alpha),$$

where  $\forall i = 1, \dots, s$   $0 \neq g_i(x) \in K[x]$  and  $\deg(g) < n$ . Since

$$1 + \sum_{i=1}^s p_i g_i(\alpha)^2 = 0,$$

it follows that

$$1 + \sum_{i=1}^s p_i g_i(x)^2 = f(x)h(x), \quad h(x) \in K[x].$$

Define  $d := \max\{\deg(g_i) : i = 1, \dots, s\}$ . Then  $d < n$  and the polynomial  $f(x)h(x)$  has degree  $2d$ . The coefficient of  $x^{2d}$  is of the form

$$\sum_{i=1}^r p_i b_i^2,$$

with  $p_i \in P$  and  $b_i \in K$ ,  $b_i \neq 0$ , so

$$\sum_{i=1}^r p_i b_i^2 >_P 0.$$

Note that  $\deg(h) = 2d - n < n$  (because  $d < n$ ) and  $2d - n$  is odd.

Let  $h_1(x)$  be an irreducible factor of  $h(x)$  of odd degree and suppose  $\beta$  is a root of  $h_1(x)$ . Then

$$\deg(h_1) = [K(\beta) : K] < [L : K] = n.$$

Since  $h_1(\beta) = 0$ , also

$$f(\beta)h(\beta) = 1 + \sum_{i=1}^s p_i g_i(\beta)^2 = 0.$$

Therefore  $\sum_{i=1}^s p_i g_i(\beta)^2 = -1 \in T_{K(\beta)}(P)$  and by Corollary 1.4  $P$  does not extend to an ordering of  $K(\beta)$ . This is in contradiction with the minimality of  $n$ .  $\square$

#### 4. REAL CLOSED FIELDS

**Definition 4.1.** (*reell abgeschlossener Körper*) A field  $K$  is said to be **real closed** if

- (1)  $K$  is real,
- (2)  $K$  has no proper real algebraic extension.

**Proposition 4.2.** (*Artin-Schreier, 1926*) Let  $K$  be a field. The following are equivalent:

- (i)  $K$  is real closed.
- (ii)  $K$  has an ordering  $P$  which does not extend to any proper algebraic extension.
- (iii)  $K$  is real, has no proper algebraic extension of odd degree, and

$$K = K^2 \cup -(K^2).$$

*Proof.* (i)  $\Rightarrow$  (ii). Trivial.

(ii)  $\Rightarrow$  (iii). Let  $P$  be an ordering which does not extend to any proper algebraic extension. By Theorem 3.1, it follows that  $K$  has no proper algebraic extension of odd degree.

Let  $b \in P$ . Then  $b = a^2$  for some  $a \in K$ , otherwise by Theorem 2.1  $P$  extends to an ordering of  $K(\sqrt{b})$ , which is a proper algebraic extension of  $K$ .

Since  $K = P \cup (-P)$  and  $P = \{a^2 : a \in K\}$ , we get (iii).

(iii)  $\Rightarrow$  (i). Note  $\text{char}(K) = 0$  and  $\sqrt{-1} \notin K$  since  $K$  is real.

Then  $K(\sqrt{-1})$  is the only proper quadratic extension of  $K$ : if  $b \in K$  but  $\sqrt{b} \notin K$  (i.e.  $b$  is not a square), then  $b = -a^2$  for some  $a \neq 0, a \in K$ , and  $K(\sqrt{b}) = K(\sqrt{-1}\sqrt{a^2}) = K(\sqrt{-1})$ .

**Claim.** Every proper algebraic extension of  $K$  contains a proper quadratic subextension.

Note that if Claim is established we are done: indeed it follows that no proper extension can be real since  $-1$  is a square in it.

Let  $L/K$  a proper algebraic extension. Without loss of generality assume that  $[L : K]$  is finite and so even. By Primitive Element Theorem we can further assume that  $L'$  is a Galois extension.

Let  $G = \text{Gal}(L'/K)$ ,  $|G| = [L' : K] = 2^a m$ ,  $a \geq 1$ ,  $m$  odd. Let  $S$  be a 2-Sylow subgroup of  $G$  (i.e.  $|S| = 2^a$ ) and let  $E := \text{Fix}(S)$ . By Galois correspondence we get:

$$[E : K] = [G : S] = m \quad \text{odd.}$$

Therefore by assumption (iii) we must have  $[E : K] = [G : S] = 1$ , so  $G = S$  is a 2-group ( $|G| = 2^a$ ) and it has a subgroup  $G_1$  of index 2. By Galois correspondence, defining  $F_1 := \text{Fix}(G_1)$  we get a quadratic subextension of  $L'/K$ .  $\square$