

POSITIVE POLYNOMIALS LECTURE NOTES

(16: 10/06/10 - BEARBEITET 14/02/19)

SALMA KUHLMANN

Contents

1. Application of Schmüdgen's Positivstellensatz to the moment problem	1
2. Schmüdgen's nichtnegativstellensatz and Hankel matrices	2
3. Finite solvability of the K -Moment Problem	3
4. Haviland's Theorem	5

1. APPLICATION OF SPSS TO THE MOMENT PROBLEM (continued)

Lemma 1.1. (Lemma 2.11 of last lecture) Let $L : \mathbb{R}[\underline{X}] \rightarrow \mathbb{R}$ be a linear functional and denote by

$$\tau : (\mathbb{Z}_+)^n \rightarrow \mathbb{R}$$

the corresponding multisequence (i.e. $\tau(\underline{k}) := L(\underline{X}^{\underline{k}}) \forall \underline{k} \in (\mathbb{Z}_+)^n$).

Fix $g \in \mathbb{R}[\underline{X}]$, $g(\underline{X}) = \sum_{\underline{k} \in \mathbb{Z}_+^n} a_{\underline{k}} \underline{X}^{\underline{k}} \in \mathbb{R}[\underline{X}]$. Then $L(h^2 g) \geq 0$ for all $h \in \mathbb{R}[\underline{X}]$ if and only if the multisequence $g(E)_\tau$ is psd.

Proof. Compute:

$$1. L(\underline{X}^{\underline{l}} g) = \sum_{\underline{k} \in \mathbb{Z}_+^n} a_{\underline{k}} \tau(\underline{k} + \underline{l}) = g(E)_\tau(\underline{l}); \text{ for all } \underline{l} \in (\mathbb{Z}_+)^n.$$

Thus if $h = \sum_i c_i \underline{X}^{k_i} \in \mathbb{R}[\underline{X}]$ then $h^2 = \sum_{i,j} c_i c_j \underline{X}^{k_i + k_j}$.

$$2. \text{ So, } L(h^2 g) = L\left[\left(\sum_{i,j} c_i c_j \underline{X}^{k_i + k_j}\right)g\right] = \sum_{i,j} c_i c_j L(\underline{X}^{k_i + k_j} g)$$

$$\stackrel{\text{[by 1.]}}{=} \sum_{i,j} g(E)_\tau(\underline{k}_i + \underline{k}_j) c_i c_j. \quad \square$$

Theorem 1.2. (Schmüdgen's NNSS) (Reformulation in terms of moment sequences) Let $K = K_S$ compact, $S = \{g_1, \dots, g_s\}$ and $\tau : (\mathbb{Z}^+)^n \rightarrow \mathbb{R}$ be a given multisequence. Then τ is a K -moment sequence if and only if the multisequences $(g_1^{e_1} \dots g_s^{e_s})(E)_\tau : (\mathbb{Z}^+)^n \rightarrow \mathbb{R}$ are all psd for all $(e_1, \dots, e_s) \in \{0, 1\}^s$. \square

Next we reformulate question (1) in 2.4 of Lecture 15 in terms of Hankel matrices:

2. SCHMÜDGEN'S NNSS AND HANKEL MATRICES

We want to understand $L(h^2g) \geq 0; h, g \in \mathbb{R}[\underline{X}]$ in terms of Hankel matrices.

Definition 2.1. A real symmetric $n \times n$ matrix A is **psd** if $\underline{x}^T A \underline{x} \geq 0 \forall \underline{x} \in \mathbb{R}^n$. An $\mathbb{N} \times \mathbb{N}$ symmetric matrix (say) A is psd if $\underline{x}^T A \underline{x} \geq 0 \forall \underline{x} \in \mathbb{R}^n$ and $\forall n \in \mathbb{N}$.

Definition 2.2. Let $L \neq 0; L : \mathbb{R}[\underline{X}] \rightarrow \mathbb{R}$ be a given linear functional. Fix $g \in \mathbb{R}[\underline{X}]$. Consider symmetric bilinear form:

$$\begin{aligned} \langle \cdot, \cdot \rangle_g : \mathbb{R}[\underline{X}] \times \mathbb{R}[\underline{X}] &\rightarrow \mathbb{R} \\ \langle h, k \rangle_g &:= L(hkg); h, k \in \mathbb{R}[\underline{X}] \end{aligned}$$

Denote by S_g the $\mathbb{N} \times \mathbb{N}$ symmetric matrix with $\alpha\beta$ -entry $\langle \underline{X}^\alpha, \underline{X}^\beta \rangle_g \forall \underline{\alpha}, \underline{\beta} \in \mathbb{N}^n$, i.e. the $\alpha\beta$ -entry of S_g is $L(\underline{X}^{\alpha+\beta} g)$.

Example. Let $g = 1$, then

$$\langle \underline{X}^\alpha, \underline{X}^\beta \rangle_1 = L(\underline{X}^{\alpha+\beta}) := s_{\underline{\alpha}+\underline{\beta}}.$$

More generally, if $g = \sum a_\gamma \underline{X}^\gamma$ then

$$\langle \underline{X}^\alpha, \underline{X}^\beta \rangle_g = L\left(\sum_\gamma a_\gamma \underline{X}^{\alpha+\beta+\gamma}\right) = \sum_\gamma a_\gamma s_{\underline{\alpha}+\underline{\beta}+\underline{\gamma}}.$$

Proposition 2.3. Let L, g be fixed as above. Then the following are equivalent:

1. $L(\sigma g) \geq 0 \forall \sigma \in \sum \mathbb{R}[\underline{X}]^2$.
2. $L(h^2g) \geq 0 \forall h \in \mathbb{R}[\underline{X}]$.
3. $\langle \cdot, \cdot \rangle_g$ is psd.
4. S_g is psd.

Proof. (1) \Leftrightarrow (2) is clear.

Since $\langle h, h \rangle_g = L(h^2g)$, (2) \Leftrightarrow (3) is clear.

(3) \Leftrightarrow (4) is also clear. \square

2.4. Example. (Hamburger) Let $n = 1$. A linear functional $L : \mathbb{R}[X] \rightarrow \mathbb{R}$ comes from a Borel measure on \mathbb{R} if and only if $L(\sigma) \geq 0$ for every $\sigma \in \sum \mathbb{R}[X]^2$.

Proof. From Haviland we know L comes from a Borel measure iff $L(f) \geq 0$ for every $f(X) \in \mathbb{R}[X], f \geq 0$ on \mathbb{R} . But $\text{Psd}(\mathbb{R}) = \sum \mathbb{R}[X]^2$ (by exercise in Real Algebraic Geometry course in WS 2009-10). So the condition is clear. \square

Remark 2.5. We can express Hamburgers's Theorem via Hankel matrix S_g with $g = 1$ the constant polynomial.

$n = 1$, so (for $i, j \in \mathbb{N}$) the ij^{th} coefficient of S_1 is $s_{i+j} = L(X^{i+j})$.

Hence, $S_1 = \begin{pmatrix} s_0 & s_1 & s_2 & \dots \\ s_1 & s_2 & \dots & \\ s_2 & \dots & \ddots & \\ \dots & \dots & & \end{pmatrix}$ is psd.

2.1. REFORMULATION OF SCHMÜDGEN'S SOLUTION TO THE MOMENT PROBLEM IN TERMS OF HANKEL MATRICES

2.6. Let $S = \{g_1, \dots, g_s\} \subseteq \mathbb{R}[X]$ and $K_S \subseteq \mathbb{R}^n$ is compact. A linear functional L on $\mathbb{R}[X]$ is represented by a Borel measure on K iff the $2^S \mathbb{N} \times \mathbb{N}$ Hankel matrices $\{S_{g_1^{e_1} \dots g_s^{e_s}} | (e_1, \dots, e_s) \in \{0, 1\}^s\}$ are psd, where $S_g := [L(X^{\underline{\alpha} + \underline{\beta}} g)]_{\underline{\alpha}, \underline{\beta}}; \underline{\alpha}, \underline{\beta} \in \mathbb{N}^n$.

3. FINITE SOLVABILITY OF THE K -MOMENT PROBLEM

Definition 3.1. Let K be a basic closed semi-algebraic subset of \mathbb{R}^n .

1. The K -moment problem (**KMP**) is **finitely solvable** if there exists S finite, $S \subseteq \mathbb{R}[X]$ such that:
 - (i) $K = K_S$, and
 - (ii) \forall linear functional L on $\mathbb{R}[X]$ we have: $L(T_S) \geq 0 \Rightarrow L(\text{Psd}(K)) \geq 0$
(equivalently, (iii) $L(T_S) \geq 0 \Rightarrow \exists \mu : L = \int d\mu$).
2. We shall say S **solves the KMP** if (i) and (ii) (equivalently (i) and (iii)) hold.

3.2. Schmüdgen's solution to the KPM for K compact b.c.s.a. Let $K \subseteq \mathbb{R}^n$ be a compact basic closed semi-algebraic set. Then S solves the KMP for any finite description S of K (i.e. for all finite $S \subseteq \mathbb{R}[X]$ with $K = K_S$).

One can restate the condition “ S solves the K -Moment problem” via the equality of two preorderings. We shall adopt this approach throughout:

Definition 3.3. Let $T_S \subseteq \mathbb{R}[\underline{X}]$ be a preordering. Define the **dual cone** of T_S :

$$T_S^v := \{L \mid L : \mathbb{R}[\underline{X}] \rightarrow \mathbb{R} \text{ is a linear functional}; L(T_S) \geq 0\},$$

and the **double dual cone**:

$$T_S^{vv} := \{f \mid f \in \mathbb{R}[\underline{X}]; L(f) \geq 0 \forall L \in T_S^v\}.$$

Lemma 3.4. For $S \subseteq \mathbb{R}[\underline{X}]$, S finite:

- (a) $T_S \subseteq T_S^{vv}$
- (b) $T_S^{vv} \subseteq \text{Psd}(K_S)$.

Proof. (a) Immediate by definition.

- (b) Let $f \in T_S^{vv}$. To show: $f(\underline{x}) \geq 0 \forall \underline{x} \in K_S$.

Now every $\underline{x} \in \mathbb{R}^n$ determines an \mathbb{R} -algebra homomorphism

$$e_{v_x} := L_{\underline{x}} \in \text{Hom}(\mathbb{R}[\underline{X}], \mathbb{R}); L_{\underline{x}}(g) = e_{v_x}(g) := g(\underline{x}) \forall g \in \mathbb{R}[\underline{X}],$$

this $L_{\underline{x}}$ is in particular a linear functional.

Moreover we claim that $L_{\underline{x}}(T_S) \geq 0$ for $\underline{x} \in K_S$. Indeed if $g \in T_S$ then $L_{\underline{x}}(g) = g(\underline{x}) \geq 0$ for $\underline{x} \in K_S$.

So, by assumption on f we must also have $L_{\underline{x}}(f) \geq 0$ for $\underline{x} \in K_S$, i.e. $f(\underline{x}) \geq 0$ for all $\underline{x} \in K_S$ as required. □

We summarize as follows:

Corollary 3.5. For finite $S \subseteq \mathbb{R}[\underline{X}]$:

$$T_S \subseteq T_S^{vv} \subseteq \text{Psd}(K_S).$$

Corollary 3.6. (Reformulation of finite solvability) Let $K \subseteq \mathbb{R}^n$ be a b.c.s.a. set and $S \subseteq \mathbb{R}[\underline{X}]$ be finite. Then S solves the KMP iff

- (j) $K = K_S$, and
- (jj) $T_S^{vv} = \text{Psd}(K)$.

Proof. Assume (ii) of definition 3.1, i.e. $\forall L : L(T_S) \geq 0 \Rightarrow L(\text{Psd}(K)) \geq 0$, and show (jj) i.e. $T_S^{\text{vv}} = \text{Psd}(K)$:

Let $f \in \text{Psd}(K)$. Show $f \in T_S^{\text{vv}}$ i.e. show $L(f) \geq 0 \forall L \in T_S^{\text{v}}$.

Assume $L(T_S) \geq 0$. Then by assumption $L(\text{Psd}(K)) \geq 0$. So, $L(f) \geq 0$ as required.

Conversely, assume (jj) and show (ii):

Let $L(T_S) \geq 0$, i.e. $L \in T_S^{\text{v}}$. Show $L(\text{Psd}(K)) \geq 0$, i.e show $L(f) \geq 0 \forall f \in \text{Psd}(K)$.

Now [by assumption (jj)] $f \in \text{Psd}(K) \Rightarrow f \in T_S^{\text{vv}} \Rightarrow L(f) \geq 0 \forall L \in T_S^{\text{v}}$. \square

We shall come back later to T_S^{vv} and describe it as closure w.r.t. an appropriate topology.

4. HAVILAND'S THEOREM

For the proof of Haviland's theorem (2.5 of lecture 15), we will recall Riesz Representation Theorem.

Definition 4.1. A topological space is said to be **Hausdorff** (or **seperated**) if it satisfies

(H4): any two distinct points have disjoint neighbourhoods, or

(T₂): two distinct points always lie in disjoint open sets.

Definition 4.2. A topological space χ is said to be **locally compact** if $\forall x \in \chi \exists$ an open neighbourhood $\mathcal{U} \ni x$ such that $\overline{\mathcal{U}}$ is compact.

Theorem 4.3. (Riesz Representation Theorem) Let χ be a locally compact Hausdorff space and $L : \text{Cont}_c(\chi, \mathbb{R}) \rightarrow \mathbb{R}$ be a positive linear functional i.e. $L(f) \geq 0 \forall f \geq 0$ on χ . Then there exists a unique (positive regular) Borel measure μ on χ such that $L(f) = \int_{\chi} f d\mu \forall f \in \text{Cont}_c(\chi, \mathbb{R})$, where $\text{Cont}_c(\chi, \mathbb{R}) :=$

the ring (\mathbb{R} -algebra) of all continuous functions $f : \chi \rightarrow \mathbb{R}$ (addition and multiplication defined pointwise) with compact support i.e. such that the set $\text{supp}(f) := \{x \in \chi : f(x) \neq 0\}$ is compact.

Definition 4.4. L **positive** means:

$$L(f) \geq 0 \forall f \in \text{Cont}_c(\chi, \mathbb{R}) \text{ with } f \geq 0 \text{ on } \chi.$$