

REAL ALGEBRAIC GEOMETRY LECTURE NOTES
(03: 20/04/15 - CORRECTED ON 02/05/2019)

SALMA KUHLMANN

CONTENTS

1.	Hahn Sandwich Proposition	1
2.	Immediate extensions	1
3.	Valuation independence	3
4.	Maximal valuation independence	4

1. HAHN SANDWICH PROPOSITION

Lemma 1.1.

(i) $\bigsqcup_{\gamma \in \Gamma} B(\gamma) \subseteq \mathbf{H}_{\gamma \in \Gamma} B(\gamma)$.

(ii)

$$\begin{aligned} S\left(\bigsqcup_{\gamma \in \Gamma} B(\gamma)\right) &\cong [\Gamma, \{B(\gamma) : \gamma \in \Gamma\}] \\ &\cong S(\mathbf{H}_{\gamma \in \Gamma} B(\gamma)). \end{aligned}$$

We shall show that if $Z = Q$ is a field and (V, v) is a valued Q -vector space with skeleton $S(V) = [\Gamma, \{B(\gamma) : \gamma \in \Gamma\}]$, then

$$\left(\bigsqcup_{\gamma \in \Gamma} B(\gamma), v_{\min}\right) \hookrightarrow (V, v) \hookrightarrow (\mathbf{H}_{\gamma \in \Gamma} B(\gamma), v_{\min}).$$

2. IMMEDIATE EXTENSIONS

Definition 2.1. Let (V_i, v_i) be valued Q -vector spaces ($i = 1, 2$).

- (1) Let $V_1 \subseteq V_2$ be a Q -subspace with $v_1(V_1) \subseteq v_2(V_2)$. We say that (V_2, v_2) is an **extension** of (V_1, v_1) , and we write

$$(V_1, v_1) \subseteq (V_2, v_2),$$

if $v_{2|_{V_1}} = v_1$.

(2) If $(V_1, v_1) \subseteq (V_2, v_2)$ and $\gamma \in v_1(V_1)$, the map

$$\begin{aligned} B_1(\gamma) &\longrightarrow B_2(\gamma) \\ x + (V_1)_\gamma &\mapsto x + (V_2)_\gamma \end{aligned}$$

is a natural identification of $B_1(\gamma)$ as a Q -subspace of $B_2(\gamma)$. The extension $(V_1, v_1) \subseteq (V_2, v_2)$ is **immediate** if $\Gamma := v_1(V_1) = v_2(V_2)$ and $\forall \gamma \in v_1(V_1)$

$$B_1(\gamma) = B_2(\gamma).$$

Equivalently, $(V_1, v_1) \subseteq (V_2, v_2)$ is immediate if $S(V_1) = S(V_2)$.

Lemma 2.2. (*Characterization of immediate extensions*)

The extension $(V_1, v_1) \subseteq (V_2, v_2)$ is immediate if and only if

$$\forall x \in V_2, x \neq 0, \exists y \in V_1 \text{ such that } v_2(x - y) > v_2(x).$$

Proof. We show that in a valued Q -vector space (V, v) , for every $x, y \in V$

$$v(x - y) > v(x) \iff \begin{cases} (i) & \gamma = v(x) = v(y) \text{ and} \\ (ii) & \pi(\gamma, x) = \pi(\gamma, y). \end{cases}$$

(\Leftarrow) Suppose (i) and (ii). So $x, y \in V^\gamma$ and $x - y \in V_\gamma$. Then $v(x - y) > \gamma = v(x)$.

(\Rightarrow) Suppose $v(x - y) > v(x)$. We show (i) and (ii).

Assume for a contradiction that $v(x) \neq v(y)$. Then $v(x - y) = \min\{v(x), v(y)\}$. So if $v(x) > v(y)$, then $v(y) = v(x - y) > v(x)$ and if $v(y) > v(x)$, then $v(x) = v(x - y) > v(x)$. Both is obviously a contradiction. Thus, $v(x) = v(y)$. (ii) is analogue. □

Example 2.3. $(\bigsqcup_{\gamma \in \Gamma} B(\gamma), v_{\min}) \subseteq (H_{\gamma \in \Gamma} B(\gamma), v_{\min})$

is an immediate extension.

Proof. Given $x \in H_{\gamma \in \Gamma} B(\gamma)$, $x \neq 0$, set

$$\gamma_0 := \min \text{support}(x) \quad \text{and} \quad x(\gamma_0) := b_0 \in B(\gamma_0).$$

Let $y \in \bigsqcup_{\gamma \in \Gamma} B(\gamma)$ such that

$$y(\gamma) = \begin{cases} 0 & \text{if } \gamma \neq \gamma_0 \\ b_0 & \text{if } \gamma = \gamma_0. \end{cases}$$

Namely $y = b_0 \chi_{\gamma_0}$, where

$$\chi_{\gamma_0}: \Gamma \longrightarrow Q$$

$$\chi_{\gamma_0}(\gamma) = \begin{cases} 1 & \text{if } \gamma = \gamma_0 \\ 0 & \text{if } \gamma \neq \gamma_0. \end{cases}$$

Then $v_{\min}(x - y) > \gamma_0 = v_{\min}(x)$ (because $(x - y)(\gamma_0) = x(\gamma_0) - y(\gamma_0) = b_0 - b_0 = 0$).

□

3. VALUATION INDEPENDENCE

Definition 3.1. $\mathcal{B} = \{x_i : i \in I\} \subseteq V \setminus \{0\}$ is Q -valuation independent if for $q_i \in Q$ with $q_i = 0$ for all but finitely many $i \in I$, we have

$$v\left(\sum_{i \in I} q_i x_i\right) = \min_{i \in I, q_i \neq 0} \{v(x_i)\}.$$

Remark 3.2.

(1) Q -linear independence $\not\Rightarrow$ Q -valuation independence.

Consider (\mathbb{Q}_2, v_{\min}) and the elements $x_1 = (1, 1)$, $x_2 = (1, 0)$.

(2) $\mathcal{B} \subseteq V \setminus \{0\}$ is Q -valuation independent $\Rightarrow \mathcal{B}$ is Q -linear independent.

Else $\exists q_i \neq 0$ with $\sum q_i x_i = 0$ and $\min\{v(x_i)\} = v(\sum q_i x_i) = \infty$, a contradiction.

Proposition 3.3. (*Characterization of valuation independence*)

Let $\mathcal{B} \subseteq V \setminus \{0\}$. Then \mathcal{B} is Q -valuation independent if and only if $\forall n \in \mathbb{N}$ and $\forall b_1, \dots, b_n \in \mathcal{B}$ pairwise distinct with $v(b_1) = \dots = v(b_n) = \gamma$, the coefficients

$$\pi(\gamma, b_1), \dots, \pi(\gamma, b_n) \in B(\gamma)$$

are Q -linear independent in the Q -vector space $B(\gamma)$.

Proof.

(\Rightarrow) Let $b_1, \dots, b_n \in \mathcal{B}$ with $v(b_1) = \dots = v(b_n) = \gamma$ and suppose for a contradiction that

$$\pi(\gamma, b_1), \dots, \pi(\gamma, b_n) \in B(\gamma)$$

are not Q -linear independent. So there are $q_1, \dots, q_n \in Q$ non-zero such that $\pi(\gamma, \sum q_i b_i) = 0$, so $v(\sum q_i b_i) > \gamma$. This contradicts the valuation independence.

(\Leftarrow) We show that

$$v\left(\sum q_i b_i\right) = \min\{v(b_i)\} = \gamma.$$

Since $\pi(\gamma, b_1), \dots, \pi(\gamma, b_n)$ are Q -linear independent in $B(\gamma)$,

$$\pi\left(\gamma, \sum q_i b_i\right) \neq 0,$$

i.e. $v(\sum q_i b_i) \leq \gamma$.

On the other hand $v(\sum q_i b_i) \geq \gamma$, so $v(\sum q_i b_i) = \gamma = \min\{v(b_i)\}$.

□

4. MAXIMAL VALUATION INDEPENDENCE

By Zorn's lemma, maximal valuation independent sets exist:

Corollary 4.1. (*Characterization of maximal valuation independent sets*)
 $\mathcal{B} \subseteq V \setminus \{0\}$ is maximal valuation independent if and only if $\forall \gamma \in v(V)$

$$\mathcal{B}_\gamma := \{\pi(\gamma, b) : b \in \mathcal{B}, v(b) = \gamma\}$$

is a Q -vector space basis of $B(\gamma)$.

Corollary 4.2. Let $\mathcal{B} \subseteq V \setminus \{0\}$ be valuation independent in (V, v) . Then \mathcal{B} is maximal valuation independent if and only if the extension

$$\langle \mathcal{B} \rangle := (V_0, v|_{V_0}) \subseteq (V, v)$$

is an immediate extension.

Proof.

(\Rightarrow) Assume $\mathcal{B} \subseteq V$ is maximal valuation independent. We show $V_0 \subseteq V$ is immediate.

If not $\exists x \in V, x \neq 0$, such that

$$\forall y \in V_0 : v(x - y) \leq v(x).$$

We will show that in this case $\mathcal{B} \cup \{x\}$ is valuation independent (which will contradict our maximality assumption). Consider $v(y_0 + qx)$, $q \in Q, q \neq 0, y_0 \in V_0$. Set $y := -y_0/q$. We claim that $v(y_0 + qx) = v(x - y) = \min\{v(x), v(y)\} = \min\{v(x), v(y_0)\}$.

This follows immediately from

Fact: $v(x - y) \leq v(x) \iff v(x - y) = \min\{v(x), v(y)\}$.

Proof of the fact. The implication (\Leftarrow) is trivial. To see (\Rightarrow), assume that $v(x - y) > \min\{v(x), v(y)\}$.

If $\min\{v(x), v(y)\} = v(x)$, then we have the contradiction

$$v(x) \geq v(x - y) > \min\{v(x), v(y)\} = v(x).$$

If $\min\{v(x), v(y)\} = v(y) < v(x)$, then $v(y) = v(x - y) > v(y)$, again a contradiction.

(\Leftarrow) Now assume that $(V_0, v|_{V_0}) \subseteq (V, v)$ is immediate. We show that \mathcal{B} is maximal valuation independent.

If not, $\mathcal{B} \cup \{x\}$ is valuation independent for some $x \in V \setminus \{0\}$ with $x \notin \mathcal{B}$. So $\forall y \in V_0$ we get $v(x - y) \leq v(x)$ by the fact above. This contradicts that $(V_0, v|_{V_0}) \subseteq (V, v)$ is immediate. □