

REAL ALGEBRAIC GEOMETRY LECTURE NOTES
(09: 17/11/2009 - BEARBEITET 22/11/2022)

SALMA KUHLMANN

CONTENTS

1.	Basic version of Tarski-Seidenberg	1
2.	Tarski Transfer Principle I	2
3.	Tarski Transfer Principle II	3
4.	Tarski Transfer Principle III	3
5.	Tarski Transfer Principle IV	4
6.	Lang's Homomorphism Theorem	4

1. BASIC VERSION OF TARSKI-SEIDENBERG

Basic version: Let (R, \leq) be a real closed field. We are interested in a system of equations and inequalities (*Gleichungen und Ungleichungen*) for $\underline{X} = (X_1, \dots, X_n)$ of the form

$$S(\underline{X}) := \begin{cases} f_1(\underline{X}) \triangleleft_1 0 \\ \vdots \\ f_k(\underline{X}) \triangleleft_k 0 \end{cases}$$

where $\forall i = 1, \dots, k \ \triangleleft_i \in \{\geq, >, =, \neq\}$ and $f_i(\underline{X}) \in \mathbb{Q}[\underline{X}]$ or $f_i(\underline{X}) \in R[\underline{X}]$. We say that $S(\underline{X})$ is a system of polynomial equalities and inequalities with coefficients in \mathbb{Q} (or with coefficients in R) in n variables.

Theorem 1.1. (*Tarski-Seidenberg Theorem: Basic Version*) *Let $S(\underline{T}; \underline{X})$ be a system with coefficients in \mathbb{Q} in $m+n$ variables, with $\underline{T} = (T_1, \dots, T_m)$ and $\underline{X} = (X_1, \dots, X_n)$. Then there exist $S_1(\underline{T}), \dots, S_l(\underline{T})$ systems in m variables and coefficients in \mathbb{Q} such that:*

for every real closed field R and every $\underline{t} = (t_1, \dots, t_m) \in R^m$ the system $S(\underline{t}; \underline{X})$ of polynomial equalities and inequalities in n variables and coefficients in R obtained by substituting T_i with t_i in $S(\underline{T}, \underline{X})$ for every $i = 1, \dots, m$, has a solution $\underline{x} = (x_1, \dots, x_n) \in R^n$ if and only if \underline{t} is a solution for one of the systems $S_1(\underline{T}), \dots, S_l(\underline{T})$.

Example 1.2. Let $m = 3$ and $n = 1$, so $\underline{T} = (T_1, T_2, T_3)$ and $\underline{X} = X$, and

$$S(\underline{T}, \underline{X}) := \left\{ T_1 X^2 + T_2 X + T_3 = 0 \right.$$

Let R be a real closed field and $(t_1, t_2, t_3) \in R^3$. Then $S(\underline{t}; X)$ has a solution in R if and only if

$$(t_1 \neq 0 \wedge t_2^2 - 4t_1t_3 \geq 0) \quad \vee \quad (t_1 = 0 \wedge t_2 \neq 0) \quad \vee \quad (t_1 = t_2 = t_3 = 0).$$

$$\begin{array}{ccc} \downarrow & & \downarrow \\ S_1(T_1, T_2, T_3) & & S_2(T_1, T_2, T_3) \end{array} \quad \vee \quad \begin{array}{ccc} \downarrow & & \downarrow \\ S_3(T_1, T_2, T_3) & & \end{array}$$

Concise version:

$$\forall \underline{T} [(\exists \underline{X} : S(\underline{T}; \underline{X})) \Leftrightarrow (\bigvee_{i=1}^l S_i(\underline{T}))].$$

Remark 1.3. The proof is by induction on n .

The case $n = 1$ is the heart of the proof and we will show it later.

For now, let us just convince ourselves that the induction step is straightforward.

Assume $n > 1$, so

$$S(\underline{T}; X_1, \dots, X_n) = S(\underline{T}, X_1, \dots, X_{n-1}; X_n).$$

By case $n = 1$ we have finitely many systems $S_1(\underline{T}, X_1, \dots, X_{n-1}), \dots, S_l(\underline{T}, X_1, \dots, X_{n-1})$ such that

for any real closed field R and any $(t_1, \dots, t_m, x_1, \dots, x_{n-1}) \in R^{m+n-1}$ we have

$$\exists X_n : S(t_1, \dots, t_m, x_1, \dots, x_{n-1}; X_n) \iff \bigvee_{i=1}^l S_i(t_1, \dots, t_m, x_1, \dots, x_{n-1}).$$

By induction hypothesis on $n - 1$:

for every fixed i , $1 \leq i \leq l$, \exists systems $S_{ij}(\underline{T})$, $j = 1, \dots, l_i$ such that: for each real closed field R and each $\underline{t} \in R^m$ the system

$$S_i(\underline{t}; X_1, \dots, X_{n-1})$$

has a solution $(x_1, \dots, x_{n-1}) \in R^{n-1}$ if and only if \underline{t} is a solution for one of the systems $S_{ij}(\underline{T})$; $j = 1, \dots, l_i$.

Therefore for any real closed field R and any $\underline{t} \in R^m$

$$S(\underline{t}; X_1, \dots, X_n) \text{ has a solution } \underline{x} \in R^n \text{ if and only if}$$

\underline{t} is a solution to one of the systems $\{S_{ij}(\underline{T}); i = 1, \dots, l, j = 1, \dots, l_i\}$

2. TARSKI TRANSFER PRINCIPLE I

Theorem 2.1. *Let $S(\underline{T}, \underline{X})$ be a system with coefficients in \mathbb{Q} in $m + n$ variables. Let (K, \leq) be an ordered field. Let R_1, R_2 be two real closed extensions of (K, \leq) . Then for every $\underline{t} \in K^m$, the system $S(\underline{t}, \underline{X})$ has a solution $\underline{x} \in R_1^n$ if and only if it has a solution $\underline{x} \in R_2^n$.*

Proof. Let $\underline{t} \in K^m \subseteq R_1^m \cap R_2^m$. There are systems $S_i(\underline{T})$ ($i = 1, \dots, l$) with coefficients in \mathbb{Q} and variables T_1, \dots, T_m such that

$$\exists \underline{x} \in R_1^n : S(\underline{t}, \underline{x}) \iff \underline{t} \text{ satisfies } \bigvee_{i=1}^l S_i(\underline{T}) \iff \exists \underline{x} \in R_2^n : S(\underline{t}, \underline{x}).$$

□

3. TARSKI TRANSFER PRINCIPLE II

Theorem 3.1. *Let (K, \leq) be an ordered field, R_1, R_2 two real closed extensions of (K, \leq) . Then a system of polynomial equations and inequalities of the form*

$$S(\underline{X}) := \begin{cases} f_1(\underline{X}) \triangleleft_1 0 \\ \vdots \\ f_k(\underline{X}) \triangleleft_k 0 \end{cases}$$

where $\forall i = 1, \dots, k \triangleleft_i \in \{\geq, >, =, \neq\}$ and $f_i(\underline{X}) \in K[X_1, \dots, X_n]$,

has a solution $\underline{x} \in R_1^n \iff$ it has a solution $\underline{x} \in R_2^n$.

Proof. Let t_1, \dots, t_m be the coefficients of the polynomials f_1, \dots, f_k , listed in some fixed order. Replacing the coefficients t_1, \dots, t_m by variables T_1, \dots, T_m yields a system $\sigma(\underline{T}, \underline{X})$ in $m + n$ variables with coefficients in \mathbb{Q} (in fact in \mathbb{Z}) for which

$$\sigma(t_1, \dots, t_m, \underline{X}) = S(\underline{X}).$$

Now we can apply Tarski Transfer I. □

4. TARSKI TRANSFER PRINCIPLE III

Theorem 4.1. *Suppose that $R \subseteq R_1$ are real closed fields. Then a system of polynomial equations and inequalities with coefficients in R*

$$S(\underline{X}) := \begin{cases} f_1(\underline{X}) \triangleleft_1 0 \\ \vdots \\ f_k(\underline{X}) \triangleleft_k 0 \end{cases}$$

where $\forall i = 1, \dots, k \triangleleft_i \in \{\geq, >, =, \neq\}$ and $f_i(\underline{X}) \in R[X_1, \dots, X_n]$

has a solution $\underline{x} \in R_1^n \iff$ it has a solution $\underline{x} \in R^n$.

Proof. Apply Tarski Transfer II with $K = R_2 = R$. □

5. TARSKI TRANSFER PRINCIPLE IV

Theorem 5.1. *Let R be a real closed field and (F, \leq) an ordered field extension of R . Then a system of polynomial equations and inequalities of the form*

$$S(\underline{X}) := \begin{cases} f_1(\underline{X}) \triangleleft_1 0 \\ \vdots \\ f_k(\underline{X}) \triangleleft_k 0 \end{cases}$$

where $\forall i = 1, \dots, k \ \triangleleft_i \in \{\geq, >, =, \neq\}$ and $f_i(\underline{X}) \in R[X_1, \dots, X_n]$

has a solution $\underline{x} \in F^n \iff$ it has a solution $\underline{x} \in R^n$.

Proof. Let R_1 be the real closure of the ordered field (F, \leq) and apply Tarski Transfer III. \square

6. LANG'S HOMOMORPHISM THEOREM

Corollary 6.1. *Suppose R and R_1 are real closed fields, $R \subseteq R_1$. Then a system of polynomial equations of the form*

$$S(\underline{X}) := \begin{cases} f_1(\underline{X}) = 0 \\ \vdots \\ f_k(\underline{X}) = 0 \end{cases} \quad f_i(\underline{x}) \in R[X_1, \dots, X_n]$$

has a solution $\underline{x} \in R_1^n$ if and only if it has a solution $\underline{x} \in R^n$.

Proof. Apply Tarski Transfer III. \square

The previous Corollary is equivalent to the following:

Theorem 6.2. (*Homomorphism Theorem I*). *Let R and R_1 be real closed fields, $R \subseteq R_1$. For any ideal $I \subseteq R[\underline{X}]$, if there exists an R -algebra homomorphism*

$$\varphi: R[\underline{X}]/I \longrightarrow R_1$$

then there exists an R -algebra homomorphism

$$\psi: R[\underline{X}]/I \longrightarrow R.$$

Proof. By Hilbert's Basis Theorem, I is finitely generated, say $I = \langle f_1, \dots, f_k \rangle$, with $f_1, \dots, f_k \in R[\underline{X}]$. Consider the system

$$S(\underline{X}) := \begin{cases} f_1(\underline{X}) = 0 \\ \vdots \\ f_k(\underline{X}) = 0 \end{cases}$$

Claim. There is a bijection

$$\{\underline{x} \in R_1^n \text{ solution to } S(\underline{X})\} \longleftrightarrow \{\varphi: R[\underline{X}]/I \rightarrow R_1 \text{ } R\text{-algebra homomorphism}\}$$

Proof of the claim:

Let $\underline{x} \in R_1^n$ be a solution to $S(\underline{X})$; then the evaluation homomorphism

$$\begin{aligned} \varphi: R[\underline{X}]/I &\longrightarrow R_1 \\ f + I &\longmapsto f(\underline{x}) \end{aligned}$$

is well-defined and is an R -algebra homomorphism.

Conversely: assume that

$$\varphi: R[\underline{X}]/I \longrightarrow R_1$$

is an R -algebra homomorphism. Then for $\underline{e} = (e_1, \dots, e_n)$ and $f = \sum \underline{a}_e \underline{X}^e = \sum a_{e_1 \dots e_n} X_1^{e_1} \dots X_n^{e_n} \in R[\underline{X}]$,

$$\varphi(f + I) = \sum \underline{a}_e \varphi(X_1 + I)^{e_1} \dots \varphi(X_n + I)^{e_n} = f(\varphi(X_1 + I), \dots, \varphi(X_n + I)).$$

In other words set $(x_1, \dots, x_n) \in R_1^n$ to be defined by $x_1 := \varphi(X_1 + I), \dots, x_n := \varphi(X_n + I)$, then (x_1, \dots, x_n) is a solution to $S(\underline{X})$ and the R -algebra homomorphism φ is indeed given by point evaluation at $\underline{x} = (x_1, \dots, x_n) \in R_1^n$.

Now apply Corollary 6.1. □