


Archimedean Ordered Fields with the Independence Property



Laura Wirth

Abstract

We specialize a conjecture on the classification of fields without the independence property to archimedean ordered fields. Specifically, we conjecture that any archimedean ordered field without the independence property is real closed. Therefore, we undertake a systematic study of the archimedean ordered fields and examine them for the independence property. We exhibit an approach for verifying the independence property and present several examples that we are considering.

Preliminaries

$\mathcal{L}_r = \{+, -, \cdot, 0, 1\}$ — language of rings

$\mathcal{L}_{or} = \{+, -, \cdot, 0, 1, <\}$ — language of ordered rings

Definition. An ordered field K is called **archimedean** if \mathbb{N} is cofinal in K .

By Hölder's Theorem any archimedean ordered field is \mathcal{L}_{or} -isomorphic to a unique subfield of \mathbb{R} .

Definition. Let \mathcal{L} be any language. An \mathcal{L} -structure \mathcal{M} has the **independence property** (in \mathcal{L}) if there exists a partitioned \mathcal{L} -formula $\varphi(\mathbf{x}; \mathbf{y})$ such that for any $n \in \mathbb{N}$, writing $[n] = \{1, \dots, n\}$, we have

$$\mathcal{M} \models \exists \mathbf{a}_1, \dots, \mathbf{a}_n \exists \mathbf{b}_0, \dots, \mathbf{b}_{[n]} : \bigwedge_{\substack{i \in [n] \\ J \subseteq [n] \\ i \in J}} \varphi(\mathbf{a}_i; \mathbf{b}_J) \wedge \bigwedge_{\substack{i \in [n] \\ J \subseteq [n] \\ i \notin J}} \neg \varphi(\mathbf{a}_i; \mathbf{b}_J).$$

Note that the independence property is preserved under elementary equivalence.

Classifying Fields without the Independence Property

The following conjecture goes back to Shelah.

Conjecture 1 [1, Introduction]. Any infinite field K that **does not have the independence property** in \mathcal{L}_r is real closed, separably closed, or admits a non-trivial henselian valuation.

Remark. Due to o-minimality a real closed field **does not have the independence property** in $\mathcal{L} \in \{\mathcal{L}_r, \mathcal{L}_{or}\}$.

Conjecture 2. Let $\mathcal{L} \in \{\mathcal{L}_r, \mathcal{L}_{or}\}$. Any (countable) subfield of \mathbb{R} that **does not have the independence property** in \mathcal{L} is real closed.

Remark. Due to the Löwenheim–Skolem Theorem we can restrict our investigations to countable subfields of \mathbb{R} .

We established the following:

Proposition 1. Conjecture 1 \Rightarrow Conjecture 2. ■

Our Approach for Verifying the Independence Property

Lemma 1. Let $\mathcal{L} \in \{\mathcal{L}_r, \mathcal{L}_{or}\}$ and let K be an (ordered) field containing an \mathcal{L} -definable subring R that has infinitely many non-associated prime elements. Then K **has the independence property** in \mathcal{L} . ■

In [2, Proposition 2] Poonen shows that \mathbb{Z} is \mathcal{L}_r -definable in any finitely generated field extension of \mathbb{Q} . We exploit this to establish the following:

Proposition 2. Let $n \in \mathbb{N}$ and let $r_1, \dots, r_n \in \mathbb{R}$. Then the field $\mathbb{Q}(r_1, \dots, r_n)$ has the independence property in \mathcal{L}_r . In particular, \mathbb{Q} and any number field in \mathbb{R} **have the independence property** in \mathcal{L}_r . ■


Acknowledgements

This work is part of my doctoral research project, which started in 2022, is supervised by Professor Salma Kuhlmann and mentored by Dr. Lothar Sebastian Krapp at Universität Konstanz.

Diagram and Examples of Archimedean Ordered Fields

We denote by

- \mathbb{Q}^{rc} the *real closure* of \mathbb{Q} , i.e. its relative algebraic closure in \mathbb{R} ,
- \mathbb{Q}^{ec} the *euclidean closure* of \mathbb{Q} ,
- \mathbb{Q}^{tr} the field $\{\alpha \in \mathbb{Q}^{rc} \mid \text{all } \mathbb{Q}\text{-conjugates of } \alpha \text{ are real}\}$ of *totally real numbers*.



The result stating that any purely transcendental extensions of \mathbb{Q} in \mathbb{R} **has the independence property** in \mathcal{L}_r is indicated on the poster of Lasse Vogel (Universität Konstanz).

Questions

1. What are prominent classes of (countable) subrings of \mathbb{R} that have infinitely many non-associated prime elements?
2. Given a (countable) field $K \subseteq \mathbb{R}$, when does K contain a subring with infinitely many non-associated prime elements that is definable in K ?
3. Let K be a proper subfield of \mathbb{Q}^{rc} . Does K have the independence property?
4. Does \mathbb{Q}^{ec} have the independence property?

References

- [1] K. Dupont, A. Hasson and S. Kuhlmann, 'Definable valuations induced by multiplicative subgroups and NIP fields', *Arch. Math. Logic* **58** (2019) 819–839.
- [2] B. Poonen, 'Uniform first-order definitions in finitely generated fields', *Duke Math. J.* **138** (2007) 1–22.

Contact

Fachbereich Mathematik und Statistik
Universität Konstanz
78457 Konstanz, Germany
laura.wirth@uni-konstanz.de
+49 7531 88 5771

