

IWOTA 2015, TBILISI, GEORGIA

Contribution by Salma Kuhlmann, Universität Konstanz, Germany

Joint work with Mehdi Ghasemi and Murray Marshall; to appear in
the Israel Journal of Mathematics

In Memoriam Murray A. Marshall: March 24 1940 - May 1st 2015
*to my colleague and friend, with my deepest gratitude for 15 years of
memorable collaboration.*

Moment problem in infinitely many variables

THE UNIVARIATE MOMENT PROBLEM

Is an old problem with origins tracing back to work of [Stieltjes](#). Given a sequence $(s_k)_{k \geq 0}$ of real numbers one wants to know when there exists a Radon measure μ on \mathbb{R} such that

$$s_k = \int x^k d\mu \quad \forall k \geq 0.^1$$

¹All Radon measures considered are assumed to be positive.

THE UNIVARIATE MOMENT PROBLEM

Is an old problem with origins tracing back to work of [Stieltjes](#). Given a sequence $(s_k)_{k \geq 0}$ of real numbers one wants to know when there exists a Radon measure μ on \mathbb{R} such that

$$s_k = \int x^k d\mu \quad \forall k \geq 0.^1$$

Since the monomials $x^k, k \geq 0$ form a basis for the polynomial algebra $\mathbb{R}[x]$, this problem is equivalent to the following one: Given a linear functional $L : \mathbb{R}[x] \rightarrow \mathbb{R}$, when does there exist a Radon measure μ on \mathbb{R} such that $L(f) = \int f d\mu \quad \forall f \in \mathbb{R}[x]$.

¹All Radon measures considered are assumed to be positive.

THE UNIVARIATE MOMENT PROBLEM

Is an old problem with origins tracing back to work of [Stieltjes](#). Given a sequence $(s_k)_{k \geq 0}$ of real numbers one wants to know when there exists a Radon measure μ on \mathbb{R} such that

$$s_k = \int x^k d\mu \quad \forall k \geq 0.^1$$

Since the monomials $x^k, k \geq 0$ form a basis for the polynomial algebra $\mathbb{R}[x]$, this problem is equivalent to the following one: Given a linear functional $L : \mathbb{R}[x] \rightarrow \mathbb{R}$, when does there exist a Radon measure μ on \mathbb{R} such that $L(f) = \int f d\mu \quad \forall f \in \mathbb{R}[x]$. One also wants to know to what extent the measure is unique, assuming it exists. [Akhiezer 1965](#) and [Shohat-Tamarkin 1943](#) are standard references.

¹All Radon measures considered are assumed to be positive.

THE MULTIVARIATE MOMENT PROBLEM

Has been considered more recently. For $n \geq 1$, $\mathbb{R}[\underline{x}] := \mathbb{R}[x_1, \dots, x_n]$ denotes the polynomial ring in n variables x_1, \dots, x_n . Given a linear functional $L : \mathbb{R}[\underline{x}] \rightarrow \mathbb{R}$ and a closed subset Y of \mathbb{R}^n one wants to know when there exists a Radon measure μ on \mathbb{R}^n supported on Y such that $L(f) = \int f d\mu \forall f \in \mathbb{R}[\underline{x}]$.

Haviland, 1936

Such a measure exists if and only if $L(\text{Pos}(Y)) \subseteq [0, \infty)$, where $\text{Pos}(Y) := \{f \in \mathbb{R}[\underline{x}] : f(x) \geq 0 \quad \forall x \in Y\}$.

THE MULTIVARIATE MOMENT PROBLEM

Has been considered more recently. For $n \geq 1$, $\mathbb{R}[\underline{x}] := \mathbb{R}[x_1, \dots, x_n]$ denotes the polynomial ring in n variables x_1, \dots, x_n . Given a linear functional $L : \mathbb{R}[\underline{x}] \rightarrow \mathbb{R}$ and a closed subset Y of \mathbb{R}^n one wants to know when there exists a Radon measure μ on \mathbb{R}^n supported on Y such that $L(f) = \int f d\mu \forall f \in \mathbb{R}[\underline{x}]$.

[Haviland, 1936](#)

Such a measure exists if and only if $L(\text{Pos}(Y)) \subseteq [0, \infty)$, where $\text{Pos}(Y) := \{f \in \mathbb{R}[\underline{x}] : f(x) \geq 0 \quad \forall x \in Y\}$.

Again, one also wants to know to what extent the measure is unique, assuming it exists. [Berg 1987](#), [Fuglede 1983](#) are general references. A major motivation here is the close connection between the multivariate moment problem and real algebraic geometry; see e.g. [Schmüdgen 1999](#), [Marshall 2008](#), [Lasserre 2013](#).

THE INFINITE-VARIATE MOMENT PROBLEM

There is work dealing with the moment problem in infinitely many variables, mainly where the linear functional in question is continuous for a certain topology.

THE INFINITE-VARIATE MOMENT PROBLEM

There is work dealing with the moment problem in infinitely many variables, mainly where the linear functional in question is continuous for a certain topology. [Albeverio-Herzberg 2008](#) applies Schmüdgen's 1999 solution of the moment problem to represent L^1 -continuous linear functionals on the vector space of polynomials of Brownian motion as integration with respect to probability measures on the Wiener space of \mathbb{R} .

THE INFINITE-VARIATE MOMENT PROBLEM

There is work dealing with the moment problem in infinitely many variables, mainly where the linear functional in question is continuous for a certain topology. [Albeverio-Herzberg 2008](#) applies Schmüdgen's 1999 solution of the moment problem to represent L^1 -continuous linear functionals on the vector space of polynomials of Brownian motion as integration with respect to probability measures on the Wiener space of \mathbb{R} .

[Berezansky-Kondratiev 1995](#), [Berezansky-Sifrin 1971](#), [Borchers-Yngvason 1975](#), [Hegerfeldt 1975](#), [Infusino-Kuna-Rota 2014](#), [Infusino 2015](#) consider continuous linear functionals on the symmetric algebra of a nuclear space.

THE INFINITE-VARIATE MOMENT PROBLEM

There is work dealing with the moment problem in infinitely many variables, mainly where the linear functional in question is continuous for a certain topology. [Albeverio-Herzberg 2008](#) applies Schmüdgen's 1999 solution of the moment problem to represent L^1 -continuous linear functionals on the vector space of polynomials of Brownian motion as integration with respect to probability measures on the Wiener space of \mathbb{R} .

[Berezansky-Kondratiev 1995](#), [Berezansky-Sifrin 1971](#), [Borchers-Yngvason 1975](#), [Hegerfeldt 1975](#), [Infusino-Kuna-Rota 2014](#), [Infusino 2015](#) consider continuous linear functionals on the symmetric algebra of a nuclear space.

[Ghasemi-Infusino-Kuhlmann-Marshall](#) (in preparation) deals with linear functionals on the symmetric algebra of a locally convex space (V, τ) which are continuous with respect to the finest locally multiplicatively convex topology extending τ .

THE INFINITE-VARIATE MOMENT PROBLEM

There is work dealing with the moment problem in infinitely many variables, mainly where the linear functional in question is continuous for a certain topology. [Albeverio-Herzberg 2008](#) applies Schmüdgen's 1999 solution of the moment problem to represent L^1 -continuous linear functionals on the vector space of polynomials of Brownian motion as integration with respect to probability measures on the Wiener space of \mathbb{R} .

[Berezansky-Kondratiev 1995](#), [Berezansky-Sifrin 1971](#), [Borchers-Yngvason 1975](#), [Hegerfeldt 1975](#), [Infusino-Kuna-Rota 2014](#), [Infusino 2015](#) consider continuous linear functionals on the symmetric algebra of a nuclear space.

[Ghasemi-Infusino-Kuhlmann-Marshall](#) (in preparation) deals with linear functionals on the symmetric algebra of a locally convex space (V, τ) which are continuous with respect to the finest locally multiplicatively convex topology extending τ . The present paper seems to be the first to deal with the general case systematically. [Today, I want to focus on the following result](#)

EXTENSION OF HAVILAND'S THEOREM

Let $A = A_\Omega := \mathbb{R}[x_i \mid i \in \Omega]$, the ring of polynomials in an arbitrary number of variables $x_i, i \in \Omega$ with coefficients in \mathbb{R} .

Extension of Haviland

Suppose $L : A_\Omega \rightarrow \mathbb{R}$ is linear and $L(\text{Pos}_{A_\Omega}(Y)) \subseteq [0, \infty)$ where Y is a closed subset of \mathbb{R}^Ω satisfying condition (i) below. Then there exists a constructibly Radon measure ν on \mathbb{R}^Ω supported by Y such that $L(f) = \int \hat{f} d\nu \forall f \in A_\Omega$.

EXTENSION OF HAVILAND'S THEOREM

Let $A = A_\Omega := \mathbb{R}[x_i \mid i \in \Omega]$, the ring of polynomials in an arbitrary number of variables $x_i, i \in \Omega$ with coefficients in \mathbb{R} .

Extension of Haviland

Suppose $L : A_\Omega \rightarrow \mathbb{R}$ is linear and $L(\text{Pos}_{A_\Omega}(Y)) \subseteq [0, \infty)$ where Y is a closed subset of \mathbb{R}^Ω satisfying condition (i) below. Then there exists a constructibly Radon measure ν on \mathbb{R}^Ω supported by Y such that $L(f) = \int \hat{f} d\nu \forall f \in A_\Omega$.

Condition (i): Y is described by countably many inequalities i.e., there exists a countable $S \subset A_\Omega$ such that $Y = \{\alpha \in \mathbb{R}^\Omega \mid \hat{g}(\alpha) \geq 0 \forall g \in S\}$.

EXTENSION OF HAVILAND'S THEOREM

Let $A = A_\Omega := \mathbb{R}[x_i \mid i \in \Omega]$, the ring of polynomials in an arbitrary number of variables $x_i, i \in \Omega$ with coefficients in \mathbb{R} .

Extension of Haviland

Suppose $L : A_\Omega \rightarrow \mathbb{R}$ is linear and $L(\text{Pos}_{A_\Omega}(Y)) \subseteq [0, \infty)$ where Y is a closed subset of \mathbb{R}^Ω satisfying condition (i) below. Then there exists a constructibly Radon measure ν on \mathbb{R}^Ω supported by Y such that $L(f) = \int \hat{f} d\nu \forall f \in A_\Omega$.

Condition (i): Y is described by countably many inequalities i.e., there exists a countable $S \subset A_\Omega$ such that $Y = \{\alpha \in \mathbb{R}^\Omega \mid \hat{g}(\alpha) \geq 0 \forall g \in S\}$. We note that Condition (i) is always satisfied for countable Ω .

EXTENSION OF HAVILAND'S THEOREM

Let $A = A_\Omega := \mathbb{R}[x_i \mid i \in \Omega]$, the ring of polynomials in an arbitrary number of variables $x_i, i \in \Omega$ with coefficients in \mathbb{R} .

Extension of Haviland

Suppose $L : A_\Omega \rightarrow \mathbb{R}$ is linear and $L(\text{Pos}_{A_\Omega}(Y)) \subseteq [0, \infty)$ where Y is a closed subset of \mathbb{R}^Ω satisfying condition (i) below. Then there exists a constructibly Radon measure ν on \mathbb{R}^Ω supported by Y such that $L(f) = \int \hat{f} d\nu \forall f \in A_\Omega$.

Condition (i): Y is described by countably many inequalities i.e., there exists a countable $S \subset A_\Omega$ such that $Y = \{\alpha \in \mathbb{R}^\Omega \mid \hat{g}(\alpha) \geq 0 \forall g \in S\}$. We note that Condition (i) is always satisfied for countable Ω .

Extension of Haviland in the countable case

Suppose Ω is countable, $L : A_\Omega \rightarrow \mathbb{R}$ is linear and $L(\text{Pos}_{A_\Omega}(Y)) \subseteq [0, \infty)$ where Y is a closed subset of \mathbb{R}^Ω . Then there exists a Radon measure ν on \mathbb{R}^Ω supported by Y such that $L(f) = \int \hat{f} d\nu \forall f \in A_\Omega$.

TERMINOLOGY, NOTATIONS, GENERAL SETTING

- ▶ All rings considered are commutative with 1.

TERMINOLOGY, NOTATIONS, GENERAL SETTING

- ▶ All rings considered are commutative with 1.
- ▶ All ring homomorphisms considered send 1 to 1.

TERMINOLOGY, NOTATIONS, GENERAL SETTING

- ▶ All rings considered are commutative with 1.
- ▶ All ring homomorphisms considered send 1 to 1.
- ▶ All rings we are interested in are \mathbb{R} -algebras.

TERMINOLOGY, NOTATIONS, GENERAL SETTING

- ▶ All rings considered are commutative with 1.
- ▶ All ring homomorphisms considered send 1 to 1.
- ▶ All rings we are interested in are \mathbb{R} -algebras.
- ▶ For a commutative ring A , $X(A)$ the **character space of A** is the set of all ring homomorphisms $\alpha : A \rightarrow \mathbb{R}$, .

TERMINOLOGY, NOTATIONS, GENERAL SETTING

- ▶ All rings considered are commutative with 1.
- ▶ All ring homomorphisms considered send 1 to 1.
- ▶ All rings we are interested in are \mathbb{R} -algebras.
- ▶ For a commutative ring A , $X(A)$ the **character space of A** is the set of all ring homomorphisms $\alpha : A \rightarrow \mathbb{R}$.
- ▶ For $a \in A$, $\hat{a} = \hat{a}_A : X(A) \rightarrow \mathbb{R}$ is defined by $\hat{a}_A(\alpha) = \alpha(a)$.

TERMINOLOGY, NOTATIONS, GENERAL SETTING

- ▶ All rings considered are commutative with 1.
- ▶ All ring homomorphisms considered send 1 to 1.
- ▶ All rings we are interested in are \mathbb{R} -algebras.
- ▶ For a commutative ring A , $X(A)$ the **character space of A** is the set of all ring homomorphisms $\alpha : A \rightarrow \mathbb{R}$.
- ▶ For $a \in A$, $\hat{a} = \hat{a}_A : X(A) \rightarrow \mathbb{R}$ is defined by $\hat{a}_A(\alpha) = \alpha(a)$.
- ▶ $X(A)$ is given the weakest topology such that the functions $\hat{a}_A, a \in A$ are continuous.

TERMINOLOGY, NOTATIONS, GENERAL SETTING

- ▶ All rings considered are commutative with 1.
- ▶ All ring homomorphisms considered send 1 to 1.
- ▶ All rings we are interested in are \mathbb{R} -algebras.
- ▶ For a commutative ring A , $X(A)$ the **character space of A** is the set of all ring homomorphisms $\alpha : A \rightarrow \mathbb{R}$, .
- ▶ For $a \in A$, $\hat{a} = \hat{a}_A : X(A) \rightarrow \mathbb{R}$ is defined by $\hat{a}_A(\alpha) = \alpha(a)$.
- ▶ $X(A)$ is given the weakest topology such that the functions $\hat{a}_A, a \in A$ are continuous.
- ▶ The only ring homomorphism from \mathbb{R} to itself is Id.

TERMINOLOGY, NOTATIONS, GENERAL SETTING

- ▶ All rings considered are commutative with 1.
- ▶ All ring homomorphisms considered send 1 to 1.
- ▶ All rings we are interested in are \mathbb{R} -algebras.
- ▶ For a commutative ring A , $X(A)$ the **character space of A** is the set of all ring homomorphisms $\alpha : A \rightarrow \mathbb{R}$, .
- ▶ For $a \in A$, $\hat{a} = \hat{a}_A : X(A) \rightarrow \mathbb{R}$ is defined by $\hat{a}_A(\alpha) = \alpha(a)$.
- ▶ $X(A)$ is given the weakest topology such that the functions $\hat{a}_A, a \in A$ are continuous.
- ▶ The only ring homomorphism from \mathbb{R} to itself is Id.
- ▶ Ring homomorphisms from $\mathbb{R}[\underline{x}]$ to \mathbb{R} correspond to point evaluations $f \mapsto f(\alpha), \alpha \in \mathbb{R}^n$. $X(\mathbb{R}[\underline{x}])$ is identified as a topological space with \mathbb{R}^n .

- ▶ A **quadratic module** of A is a subset M of A satisfying

$$1 \in M, M + M \subseteq M \text{ and } a^2M \subseteq M \text{ for each } a \in A.$$

- ▶ A **quadratic preordering** of A is a quadratic module of A which is also closed under multiplication.

- ▶ A **quadratic module** of A is a subset M of A satisfying

$$1 \in M, M + M \subseteq M \text{ and } a^2M \subseteq M \text{ for each } a \in A.$$

- ▶ A **quadratic preordering** of A is a quadratic module of A which is also closed under multiplication.
- ▶ For a subset X of $X(A)$,

$$\text{Pos}_A(X) := \{a \in A \mid \hat{a}_A \geq 0 \text{ on } X\}$$

is a preordering of A .

- ▶ A **quadratic module** of A is a subset M of A satisfying

$$1 \in M, M + M \subseteq M \text{ and } a^2M \subseteq M \text{ for each } a \in A.$$

- ▶ A **quadratic preordering** of A is a quadratic module of A which is also closed under multiplication.
- ▶ For a subset X of $X(A)$,

$$\text{Pos}_A(X) := \{a \in A \mid \hat{a}_A \geq 0 \text{ on } X\}$$

is a preordering of A .

- ▶ $\sum A^2$ the set of all finite sums $\sum a_i^2, a_i \in A$. It is the unique smallest quadratic module (preordering) of A .

- ▶ A **quadratic module** of A is a subset M of A satisfying

$$1 \in M, M + M \subseteq M \text{ and } a^2M \subseteq M \text{ for each } a \in A.$$

- ▶ A **quadratic preordering** of A is a quadratic module of A which is also closed under multiplication.
- ▶ For a subset X of $X(A)$,

$$\text{Pos}_A(X) := \{a \in A \mid \hat{a}_A \geq 0 \text{ on } X\}$$

is a preordering of A .

- ▶ $\sum A^2$ the set of all finite sums $\sum a_i^2, a_i \in A$. It is the unique smallest quadratic module (preordering) of A .
- ▶ For a subset $S \subseteq A$,

$$X_S := \{\alpha \in X(A) \mid \hat{a}_A(\alpha) \geq 0 \forall a \in S\}.$$

- ▶ A quadratic module M in A is **archimedean** if for each $a \in A$ there exists an integer k such that $k \pm a \in M$.

- ▶ A quadratic module M in A is **archimedean** if for each $a \in A$ there exists an integer k such that $k \pm a \in M$.
- ▶ If M is a quadratic module of A which is archimedean then X_M is compact.

- ▶ A quadratic module M in A is **archimedean** if for each $a \in A$ there exists an integer k such that $k \pm a \in M$.
- ▶ If M is a quadratic module of A which is archimedean then X_M is compact.

Archimedean Positivstellensatz

Suppose M is an archimedean quadratic module of A . Then, for any $a \in A$, the following are equivalent:

- (1) $\hat{a}_A \geq 0$ on X_M .
- (2) $a + \epsilon \in M$ for all real $\epsilon > 0$.

CONSTRUCTIBLY BOREL SETS

- ▶ The open sets

$$U_A(a) := \{\alpha \in X(A) \mid \hat{a}_A(\alpha) > 0\}, \quad a \in A$$

form a basis for the topology on $X(A)$

CONSTRUCTIBLY BOREL SETS

- ▶ The open sets

$$U_A(a) := \{\alpha \in X(A) \mid \hat{a}_A(\alpha) > 0\}, \quad a \in A$$

form a basis for the topology on $X(A)$

- ▶ If A is generated as an \mathbb{R} -algebra by $x_i, i \in \Omega$, the embedding $X(A) \hookrightarrow \mathbb{R}^\Omega$ defined by $\alpha \mapsto (\alpha(x_i))_{i \in \Omega}$ identifies $X(A)$ with a subspace of \mathbb{R}^Ω .

CONSTRUCTIBLY BOREL SETS

- ▶ The open sets

$$U_A(a) := \{\alpha \in X(A) \mid \hat{a}_A(\alpha) > 0\}, \quad a \in A$$

form a basis for the topology on $X(A)$

- ▶ If A is generated as an \mathbb{R} -algebra by $x_i, i \in \Omega$, the embedding $X(A) \hookrightarrow \mathbb{R}^\Omega$ defined by $\alpha \mapsto (\alpha(x_i))_{i \in \Omega}$ identifies $X(A)$ with a subspace of \mathbb{R}^Ω .
- ▶ Sets of the form

$$\{b \in \mathbb{R}^\Omega \mid \sum_{i \in I} (b_i - p_i)^2 < r\},$$

where $r, p_i \in \mathbb{Q}$ and I is a finite subset of Ω , form a basis for the product topology on \mathbb{R}^Ω .

- ▶ It follows that sets of the form

$$U_A(r - \sum_{i \in I} (x_i - p_i)^2), \quad r, p_i \in \mathbb{Q}, \quad I \text{ a finite subset of } \Omega, \quad (1)$$

form a basis for $X(A)$.

- ▶ It follows that sets of the form

$$U_A(r - \sum_{i \in I} (x_i - p_i)^2), \quad r, p_i \in \mathbb{Q}, \quad I \text{ a finite subset of } \Omega, \quad (1)$$

form a basis for $X(A)$.

- ▶ A subset E of $X(A)$ is called **Borel** if E is an element of the σ -algebra of subsets of $X(A)$ generated by the open sets.

- ▶ It follows that sets of the form

$$U_A(r - \sum_{i \in I} (x_i - p_i)^2), \quad r, p_i \in \mathbb{Q}, \quad I \text{ a finite subset of } \Omega, \quad (1)$$

form a basis for $X(A)$.

- ▶ A subset E of $X(A)$ is called **Borel** if E is an element of the σ -algebra of subsets of $X(A)$ generated by the open sets.
- ▶ A subset E of $X(A)$ is said to be **constructible** (resp., **constructibly Borel**) if E is an element of the algebra (resp., σ -algebra) of subsets of $X(A)$ generated by $U_A(a)$, $a \in A$.
- ▶ Clearly Constructible \Rightarrow constructibly Borel \Rightarrow Borel.

Countably generated algebras

If A is generated as an \mathbb{R} -algebra by a countable set $\{x_i \mid i \in \Omega\}$ then every Borel set of $X(A)$ is constructibly Borel.

Proof.

Sets of the form (1) form a countable basis for the topology on $X(A)$. □

SUPPORT

- ▶ The support of a measure is not defined in general. For a measure space (X, Σ, μ) and a subset Y of X , we say μ is supported by Y if $E \cap Y = \emptyset \Rightarrow \mu(E) = 0 \forall E \in \Sigma$.

SUPPORT

- ▶ The support of a measure is not defined in general. For a measure space (X, Σ, μ) and a subset Y of X , we say μ is supported by Y if $E \cap Y = \emptyset \Rightarrow \mu(E) = 0 \forall E \in \Sigma$.
- ▶ In this situation, if $\Sigma' := \{E \cap Y \mid E \in \Sigma\}$, and $\mu'(E \cap Y) := \mu(E) \forall E \in \Sigma$, then Σ' is a σ -algebra of subsets of Y , μ' is a well-defined measure on (Y, Σ') , the inclusion map $i : Y \rightarrow X$ is a measurable function, and μ is the pushforward of μ' to X .

SUPPORT

- ▶ The support of a measure is not defined in general. For a measure space (X, Σ, μ) and a subset Y of X , we say μ is supported by Y if $E \cap Y = \emptyset \Rightarrow \mu(E) = 0 \forall E \in \Sigma$.
- ▶ In this situation, if $\Sigma' := \{E \cap Y \mid E \in \Sigma\}$, and $\mu'(E \cap Y) := \mu(E) \forall E \in \Sigma$, then Σ' is a σ -algebra of subsets of Y , μ' is a well-defined measure on (Y, Σ') , the inclusion map $i : Y \rightarrow X$ is a measurable function, and μ is the pushforward of μ' to X .
- ▶ If (Y, Σ', μ') is a measure space, (X, Σ) is a σ -algebra, $i : Y \rightarrow X$ is any measurable function, and μ is the pushforward of μ' to (X, Σ) , then for each measurable function $f : X \rightarrow \mathbb{R}$, $\int f d\mu = \int (f \circ i) d\mu'$ (change in variables theorem).

CONSTRUCTIBLY RADON MEASURES

- ▶ A **Radon measure** on $X(A)$ is a positive measure μ on the σ -algebra of Borel sets of $X(A)$ which is locally finite (every point has a neighbourhood of finite measure) and inner regular (each Borel set can be approximated from within using a compact set).

CONSTRUCTIBLY RADON MEASURES

- ▶ A **Radon measure** on $X(A)$ is a positive measure μ on the σ -algebra of Borel sets of $X(A)$ which is locally finite (every point has a neighbourhood of finite measure) and inner regular (each Borel set can be approximated from within using a compact set).
- ▶ A **constructibly Radon measure** on $X(A)$ is a positive measure μ on the σ -algebra of constructibly Borel sets of $X(A)$ such that for, each countably generated subalgebra A' of A , the pushforward of μ to $X(A')$ via the restriction map $\alpha \mapsto \alpha|_{A'}$ is a Radon measure on $X(A')$.

From now on we consider only Radon and constructibly Radon measures having the additional property that \hat{a}_A is μ -integrable (i.e., $\int \hat{a}_A d\mu$ is well-defined and finite) for all $a \in A$.

THE MOMENT PROBLEM IN THIS GENERAL SETTING

- ▶ For a linear functional $L : A \rightarrow \mathbb{R}$, one can consider the set of Radon or constructibly Radon measures μ on $X(A)$ such that $L(a) = \int \hat{a}_A d\mu \forall a \in A$. The **moment problem** is to understand this set of measures, for a given linear functional $L : A \rightarrow \mathbb{R}$. In particular, one wants to know: (i) When is this set non-empty? (ii) In case it is non-empty, when is it a singleton set?

THE MOMENT PROBLEM IN THIS GENERAL SETTING

- ▶ For a linear functional $L : A \rightarrow \mathbb{R}$, one can consider the set of Radon or constructibly Radon measures μ on $X(A)$ such that $L(a) = \int \hat{a}_A d\mu \forall a \in A$. The **moment problem** is to understand this set of measures, for a given linear functional $L : A \rightarrow \mathbb{R}$. In particular, one wants to know: (i) When is this set non-empty? (ii) In case it is non-empty, when is it a singleton set?
- ▶ A linear functional $L : A \rightarrow \mathbb{R}$ is said to be **positive** if $L(\sum A^2) \subseteq [0, \infty)$ and **M -positive** for some quadratic module M of A , if $L(M) \subseteq [0, \infty)$.

TWO SPECIAL ALGEBRAS; TOWARDS THE PROOF OF THE MAIN RESULT

Let Ω is an arbitrary index set.

- ▶ As above, $A = A_\Omega := \mathbb{R}[x_i \mid i \in \Omega]$, we further define

TWO SPECIAL ALGEBRAS; TOWARDS THE PROOF OF THE MAIN RESULT

Let Ω is an arbitrary index set.

- ▶ As above, $A = A_\Omega := \mathbb{R}[x_i \mid i \in \Omega]$, we further define
- ▶ $B = B_\Omega := \mathbb{R}[x_i, \frac{1}{1+x_i^2} \mid i \in \Omega]$, the localization of A at the multiplicative set generated by the $1 + x_i^2, i \in \Omega$, and

TWO SPECIAL ALGEBRAS; TOWARDS THE PROOF OF THE MAIN RESULT

Let Ω is an arbitrary index set.

- ▶ As above, $A = A_\Omega := \mathbb{R}[x_i \mid i \in \Omega]$, we further define
- ▶ $B = B_\Omega := \mathbb{R}[x_i, \frac{1}{1+x_i^2} \mid i \in \Omega]$, the localization of A at the multiplicative set generated by the $1 + x_i^2, i \in \Omega$, and
- ▶ $C = C_\Omega := \mathbb{R}[\frac{1}{1+x_i^2}, \frac{x_i}{1+x_i^2} \mid i \in \Omega]$, the \mathbb{R} -subalgebra of B generated by the elements $\frac{1}{1+x_i^2}, \frac{x_i}{1+x_i^2}, i \in \Omega$.

TWO SPECIAL ALGEBRAS; TOWARDS THE PROOF OF THE MAIN RESULT

Let Ω is an arbitrary index set.

- ▶ As above, $A = A_\Omega := \mathbb{R}[x_i \mid i \in \Omega]$, we further define
- ▶ $B = B_\Omega := \mathbb{R}[x_i, \frac{1}{1+x_i^2} \mid i \in \Omega]$, the localization of A at the multiplicative set generated by the $1 + x_i^2, i \in \Omega$, and
- ▶ $C = C_\Omega := \mathbb{R}[\frac{1}{1+x_i^2}, \frac{x_i}{1+x_i^2} \mid i \in \Omega]$, the \mathbb{R} -subalgebra of B generated by the elements $\frac{1}{1+x_i^2}, \frac{x_i}{1+x_i^2}, i \in \Omega$.
- ▶ Elements of $X(A)$ and $X(B)$ are naturally identified with point evaluations $f \mapsto f(\alpha), \alpha \in \mathbb{R}^\Omega$.

TWO SPECIAL ALGEBRAS; TOWARDS THE PROOF OF THE MAIN RESULT

Let Ω is an arbitrary index set.

- ▶ As above, $A = A_\Omega := \mathbb{R}[x_i \mid i \in \Omega]$, we further define
- ▶ $B = B_\Omega := \mathbb{R}[x_i, \frac{1}{1+x_i^2} \mid i \in \Omega]$, the localization of A at the multiplicative set generated by the $1 + x_i^2, i \in \Omega$, and
- ▶ $C = C_\Omega := \mathbb{R}[\frac{1}{1+x_i^2}, \frac{x_i}{1+x_i^2} \mid i \in \Omega]$, the \mathbb{R} -subalgebra of B generated by the elements $\frac{1}{1+x_i^2}, \frac{x_i}{1+x_i^2}, i \in \Omega$.
- ▶ Elements of $X(A)$ and $X(B)$ are naturally identified with point evaluations $f \mapsto f(\alpha), \alpha \in \mathbb{R}^\Omega$.
- ▶ $X(A) = X(B) = \mathbb{R}^\Omega$, not just as sets, but also as topological spaces, giving \mathbb{R}^Ω the product topology.

We show how the moment problem for A_Ω reduces to understanding the extensions of a linear functional $L : A_\Omega \rightarrow \mathbb{R}$ to a positive linear functional on B_Ω and prove that positive linear functionals $L : B_\Omega \rightarrow \mathbb{R}$ correspond bijectively to constructibly Radon measures on \mathbb{R}^Ω .

Results in Marshall 2003

By definition, A (resp., B , resp., C) is the direct limit of the \mathbb{R} -algebras A_I (resp., B_I , resp., C_I), I running through all finite subsets of Ω . Because of this, many questions about A , B and C reduce immediately to the case where Ω is finite.

We show how the moment problem for A_Ω reduces to understanding the extensions of a linear functional $L : A_\Omega \rightarrow \mathbb{R}$ to a positive linear functional on B_Ω and prove that positive linear functionals $L : B_\Omega \rightarrow \mathbb{R}$ correspond bijectively to constructibly Radon measures on \mathbb{R}^Ω .

Results in Marshall 2003

By definition, A (resp., B , resp., C) is the direct limit of the \mathbb{R} -algebras A_I (resp., B_I , resp., C_I), I running through all finite subsets of Ω . Because of this, many questions about A , B and C reduce immediately to the case where Ω is finite.

These algebras were studied extensively in Marshall 2003 for finite Ω .

THE ARCHITECTURE OF THE PROOF

The following is a simple modification of the argument in Marshall 2003 for arbitrary Ω

THE ARCHITECTURE OF THE PROOF

The following is a simple modification of the argument in Marshall 2003 for arbitrary Ω

Extendibility from A to B

Suppose $L : A \rightarrow \mathbb{R}$ is an $\text{Pos}_A(Y)$ -positive linear functional for some closed set $Y \subseteq \mathbb{R}^\Omega$. Then L extends to an $\text{Pos}_B(Y)$ -positive linear functional $L : B \rightarrow \mathbb{R}$.

THE ARCHITECTURE OF THE PROOF

The following is a simple modification of the argument in Marshall 2003 for arbitrary Ω

Extendibility from A to B

Suppose $L : A \rightarrow \mathbb{R}$ is an $\text{Pos}_A(Y)$ -positive linear functional for some closed set $Y \subseteq \mathbb{R}^\Omega$. Then L extends to an $\text{Pos}_B(Y)$ -positive linear functional $L : B \rightarrow \mathbb{R}$.

Positive functionals on C ; Marshall 2003

Positive linear functionals $L : B \rightarrow \mathbb{R}$ restrict to positive linear functionals on C . **The cone of sums of squares of C is archimedean.** Positive linear functionals $L : C \rightarrow \mathbb{R}$ are in natural one-to-one correspondence with Radon measures μ on the compact space $X(C)$ via $L \leftrightarrow \mu$ iff $L(f) = \int \hat{f}_C d\mu \forall f \in C$.

Main Lemma

For each positive linear functional $L : B \rightarrow \mathbb{R}$ there exists a unique Radon measure μ on $X(C)$ such that $L(f) = \int \hat{f}_C d\mu \forall f \in C$. This satisfies $\mu(\Delta_i) = 0 \forall i \in \Omega$ and $L(f) = \int \tilde{f} d\mu \forall f \in B$.

Main Lemma

For each positive linear functional $L : B \rightarrow \mathbb{R}$ there exists a unique Radon measure μ on $X(C)$ such that $L(f) = \int \hat{f}_C d\mu \forall f \in C$. This satisfies $\mu(\Delta_i) = 0 \forall i \in \Omega$ and $L(f) = \int \tilde{f} d\mu \forall f \in B$.

Positive functionals on B

There is a canonical one-to-one correspondence $L \leftrightarrow \nu$ given by $L(f) = \int \hat{f}_B d\nu \forall f \in B$ between positive linear functionals L on B and constructibly Radon measures ν on $X(B)$.

The proof of the main theorem then proceeds as follows: Given L , there exists an extension of L to a linear functional L on B_Ω such that $L(\text{Pos}_{B_\Omega}(Y)) \subseteq [0, \infty)$. Denote by ν the constructibly Radon measure on \mathbb{R}^Ω corresponding to this extension. Fix a countable set S in A_Ω such that $Y = X_S$. For each $g \in S$, choose $g' \in C_\Omega$ of the form $g' = g/p_g$ for some suitably chosen element $p_g = (1 + x_{j_1}^2)^{e_1} \dots (1 + x_{j_k}^2)^{e_k}$. Let $S' = \{g' \mid g \in S\}$. Let Q' = the quadratic module of C_Ω generated by S' , Q = the quadratic module of B_Ω generated by S . Note that Q is also the quadratic module in B_Ω generated by S' , and $Q' \subseteq Q \subseteq \text{Pos}_{B_\Omega}(Y)$, so $L'(Q') \subseteq [0, \infty)$ where $L' := L|_{C_\Omega}$. By [Marshall 2003](#) there exists a Radon measure μ on $X(C_\Omega)$ supported by $X_{Q'}$ such that $L'(f) = \int \hat{f} d\mu \forall f \in C_\Omega$. Uniqueness implies that μ is the Radon measure on $X(C_\Omega)$ defined in [Main Lemma](#). One checks that ν is supported by $X_{Q'} \cap X(B_\Omega) = X_Q = X_S = Y$