

# POSITIVE POLYNOMIALS LECTURE NOTES

## (15: 08/06/10 - BEARBEITET 12/02/19)

SALMA KUHLMANN

### Contents

1. Schmüdgen's Nichtnegativstellensatz 1
2. Application of Schmüdgen's Positivstellensatz to the moment problem 2

#### 1. SCHMÜDGEN'S NICHTNEGATIVSTELLENSATZ

**1.1. Schmüdgen's Nichtnegativstellensatz** (Recall 2.2.1 of lecture 14): Let  $K_S$  be a compact basic closed semi algebraic set and  $f \in \mathbb{R}[X]$ . Then

$$f \geq 0 \text{ on } K_S \Rightarrow \forall \epsilon \text{ real, } \epsilon > 0 : f + \epsilon \in T_S.$$

**Corollary 1.2.** Let  $K = K_S$  be a compact basic closed semi algebraic set and  $L \neq 0$  be a linear functional  $L : \mathbb{R}[X] \rightarrow \mathbb{R}$  with  $L(r) = r \forall r \in \mathbb{R}$ . Then

$$\underbrace{L(T_S) \geq 0}_{\text{(i.e. } L(f) \geq 0 \forall f \in T_S)} \Rightarrow \underbrace{L(\text{Psd}(K_S)) \geq 0}_{\text{(i.e. } L(f) \geq 0 \forall f \geq 0 \text{ on } K_S)}.$$

*Proof.* W.l.o.g.  $L(1) = 1, L \neq 0$ . Let  $f \in \text{Psd}(K_S)$  and assume  $L(T_S) \geq 0$ ,

To show:  $L(f) \geq 0$

By 1.1,  $\forall \epsilon > 0 : f + \epsilon \in T_S$

So,  $L(f + \epsilon) \geq 0$  i.e.  $L(f) \geq -\epsilon \forall \epsilon > 0$  real

$\Rightarrow L(f) \geq 0$ . □

We shall now relate this to the problem of representation of linear functionals via integration along measures (i.e.  $\int d\mu$ ).

## 2. APPLICATION OF SPSS TO THE MOMENT PROBLEM

Let  $\mathcal{X}$  be a Hausdorff topological space.

**Definition 2.1.**  $\mathcal{X}$  is **locally compact** if  $\forall x \in \mathcal{X} \exists$  open  $\mathcal{U}$  in  $\mathcal{X}$  s.t.  $x \in \mathcal{U}$  and  $\overline{\mathcal{U}}$  (closure) is compact.

**Notation 2.2.**  $\mathcal{B}^\delta(\mathcal{X}) :=$  set of Borel measurable sets in  $\mathcal{X}$   
 = the smallest family of subsets of  $\mathcal{X}$  containing all compact subsets of  $\mathcal{X}$ , closed under finite  $\cup$ , set theoretic difference  $A \setminus B$  and countable  $\cap$ .

**Definition 2.3.** A **Borel measure**  $\mu$  on  $\mathcal{X}$  is a positive measure on  $\mathcal{X}$  s.t. every set in  $\mathcal{B}^\delta(\mathcal{X})$  is measurable. We also require our measure to be **regular** i.e.  $\forall B \in \mathcal{B}^\delta(\mathcal{X})$  and  $\forall \epsilon > 0 \exists K, \mathcal{U} \in \mathcal{B}^\delta(\mathcal{X}), K$  compact,  $\mathcal{U}$  open s.t.  $K \subseteq B \subseteq \mathcal{U}$  and  $\mu(K) + \epsilon \geq \mu(B) \geq \mu(\mathcal{U}) - \epsilon$ .

**2.4. Moment problem** is the following:

Given a closed set  $K \subseteq \mathbb{R}^n$  and a linear functional  $L : \mathbb{R}[\underline{X}] \rightarrow \mathbb{R}$

Question:

$$\text{when does } \exists \text{ a Borel measure } \mu \text{ on } K \text{ s.t. } \forall f \in \mathbb{R}[\underline{X}] : L(f) = \int f d\mu ? \quad (1)$$

$$\text{Necessary condition for (1): } \forall f \in \mathbb{R}[\underline{X}], f \geq 0 \text{ on } K \Rightarrow L(f) \geq 0 \quad (2)$$

$$\text{in other words: } L(\text{Psd}(K)) \geq 0 \quad (3)$$

Is this necessary condition also sufficient?

The answer is YES.

**Theorem 2.5.** (Haviland) Given  $K \subseteq \mathbb{R}^n$  closed and  $L : \mathbb{R}[\underline{X}] \rightarrow \mathbb{R}$  a linear functional with  $L(1) > 0$ :

$$\exists \mu \text{ as in (1) iff } \forall f \in \mathbb{R}[\underline{X}] : L(f) \geq 0 \text{ if } f \geq 0 \text{ on } K.$$

We shall prove Haviland's Theorem later. For now we shall deduce a corollary to SPSS.

**Corollary 2.6.** Let  $K_S = \{x \mid g_i(x) \geq 0; i = 1, \dots, s\} \subseteq \mathbb{R}^n$  be a basic closed semi-algebraic set and compact,  $L : \mathbb{R}[\underline{X}] \rightarrow \mathbb{R}$  a linear functional with  $L(1) > 0$ . If

$$L(T_S) \geq 0, \text{ then } \exists \mu \text{ positive Borel measure on } K \text{ s.t. } L(f) = \int_{K_S} f d\mu \quad \forall f \in \mathbb{R}[\underline{X}].$$

**Remark 2.7.** Let  $S = \{g_1, \dots, g_s\}$ .

1.  $L(T_S) \geq 0$  can be written as

$$L(h^2 g_1^{e_1} \dots g_s^{e_s}) \geq 0 \quad \forall h \in \mathbb{R}[\underline{X}], e_1, \dots, e_s \in \{0, 1\}.$$

2. Compare Haviland to Schmüdgen's moment problem, for compact  $K_S$ : we do not need to check  $L(\text{Psd}(K_S)) \geq 0$  we only need to check  $L(T_S) \geq 0$ .

3. Reformulation of question (1) (in 2.4) in terms of moment sequences:

Let  $L : \mathbb{R}[\underline{X}] \rightarrow \mathbb{R}$ , with  $L(1) = 1$ . Consider  $\{X^\alpha = X_1^{\alpha_1} \dots X_n^{\alpha_n}; \alpha \in \mathbb{N}_0^n\}$  a monomial basis for  $\mathbb{R}[\underline{X}]$ . So  $L$  is completely determined by the (multi)sequence of real numbers  $\tau(\alpha) := L(X^\alpha)$ ;  $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}_0^n$  (i.e.  $\tau : \mathbb{N}_0^n \rightarrow \mathbb{R}$  is a function) and conversely, every such sequence determines a linear functional  $L$ :

$$L\left(\sum_{\alpha} a_{\alpha} X^{\alpha}\right) := \sum_{\alpha} a_{\alpha} L(X^{\alpha}).$$

So, (1) (in 2.4) can be reformulated as:

Given  $K \subseteq \mathbb{R}^n$  closed, and a multisequence  $\tau = \tau(\alpha)_{\alpha \in \mathbb{N}_0^n}$  of real numbers,  $\exists \mu$  positive borel measure on  $K$  s.t  $\int X^\alpha d\mu = \tau_\alpha$  for all  $\alpha \in \mathbb{N}_0^n$ ?

**Definition 2.8.** A function  $\tau : \mathbb{N}_0^n \rightarrow \mathbb{R}$  is a  $K$ -**moment sequence** if  $\exists \mu$  positive borel measure on  $K$  s.t  $\tau(\alpha) = \int_K X^\alpha d\mu$  for all  $\alpha \in \mathbb{N}_0^n$

So (1) can be reformulated as: given  $K$  and a function  $\tau : \mathbb{N}_0^n \rightarrow \mathbb{R}$ , when is  $\tau$  a  $K$ -moment sequence?

**Definition 2.9.** A function  $\tau : (\mathbb{Z}_+)^n \rightarrow \mathbb{R}$  is called **psd** if

$$\sum_{i,j=1}^m \tau(\underline{k}_i + \underline{k}_j) c_i c_j \geq 0,$$

for  $m \geq 1$ , arbitrary distinct  $\underline{k}_1, \dots, \underline{k}_m \in (\mathbb{Z}_+)^n$ ;  $c_1, \dots, c_m \in \mathbb{R}$ .

**Definition 2.10.** Given  $\tau : (\mathbb{Z}_+)^n \rightarrow \mathbb{R}$  a function and a fixed polynomial

$g(\underline{X}) = \sum_{\underline{k} \in \mathbb{Z}_+^n} a_{\underline{k}} X^{\underline{k}} \in \mathbb{R}[\underline{X}]$ . Define a new function  $g(E)_\tau : (\mathbb{Z}_+)^n \rightarrow \mathbb{R}$  by

$$g(E)_\tau(\underline{l}) := \sum_{\underline{k} \in \mathbb{Z}_+^n} a_{\underline{k}} \tau(\underline{k} + \underline{l}); \text{ for any } \underline{l} \in (\mathbb{Z}_+)^n.$$

**Lemma 2.11.** Let  $L : \mathbb{R}[\underline{X}] \rightarrow \mathbb{R}$  be a linear functional and denote by

$$\tau : (\mathbb{Z}_+)^n \rightarrow \mathbb{R}$$

the corresponding multisequence (i.e.  $\tau(\underline{k}) := L(\underline{X}^{\underline{k}}) \forall \underline{k} \in (\mathbb{Z}_+)^n$ ).

Fix  $g \in \mathbb{R}[\underline{X}]$ . Then  $L(h^2 g) \geq 0$  for all  $h \in \mathbb{R}[\underline{X}]$  if and only if the multisequence  $g(E)_\tau$  is psd.